

# Anti-Human BCA-1 Antibody

Catalog # ABG10021

## Product Information

---

|                    |            |
|--------------------|------------|
| <b>Application</b> | WB, E      |
| <b>Reactivity</b>  | Human      |
| <b>Host</b>        | Rabbit     |
| <b>Clonality</b>   | Polyclonal |

## Additional Information

---

|                       |                                                                                                                                                                                                                                                                                                                                                             |
|-----------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Preparation</b>    | Produced from sera of rabbits pre-immunized with highly pure (>98%) recombinant hBCA-1. Anti-Human BCA-1 specific antibody was purified by affinity chromatography employing immobilized hBCA-1 matrix.                                                                                                                                                     |
| <b>WesternBlot</b>    | To detect hBCA-1 by Western Blot analysis this antibody can be used at a concentration of 0.1-0.2 µg/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant hBCA-1 is 1.5-3.0 ng/lane, under either reducing or non-reducing conditions.                                                                            |
| <b>Sandwich</b>       | To detect hBCA-1 by sandwich ELISA (using 100 µl/well antibody solution) a concentration of 0.5 - 2.0 µg/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with BioGems' Biotinylated Anti-Human BCA-1 (60-072BT) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant hBCA-1. |
| <b>Formulation</b>    | A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.                                                                                                                                                                                                                                                                                      |
| <b>Reconstitution</b> | Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.                                                                                                                                                                                                                                                        |
| <b>Storage</b>        | -20°C                                                                                                                                                                                                                                                                                                                                                       |
| <b>Precautions</b>    | Anti-Human BCA-1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.                                                                                                                                                                                                                                                 |

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.