

# Anti-Human Betacellulin Antibody

Catalog # ABG10033

## Product Information

---

|                    |            |
|--------------------|------------|
| <b>Application</b> | WB, IHC, E |
| <b>Reactivity</b>  | Human      |
| <b>Host</b>        | Rabbit     |
| <b>Clonality</b>   | Polyclonal |

## Additional Information

---

|                             |                                                                                                                                                                                                                                                                                                                                                                                                                     |
|-----------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Preparation</b>          | Produced from sera of rabbits pre-immunized with highly pure (>98%) recombinant hBetacellulin. Anti-Human Betacellulin specific antibody was purified by affinity chromatography employing immobilized hBetacellulin matrix.                                                                                                                                                                                        |
| <b>WesternBlot</b>          | To detect hBetacellulin by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 µg/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant hBetacellulin is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.                                                                                                                  |
| <b>Sandwich</b>             | To detect hBetacellulin by sandwich ELISA (using 100 µl/well antibody solution) a concentration of 0.5 - 2.0 µg/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with BioGems' Biotinylated Anti-Human Betacellulin (60-081BT) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant hBetacellulin.                                    |
| <b>Immunohistochemistry</b> | This antibody stained formalin-fixed paraffin-embedded sections of human pancreas adenocarcinoma tissue. The recommended concentration is 2.5 µg/ml with an overnight incubation at 4°C. An HRP-labeled polymer detection system was used with an alcohol-soluble AEC chromogen. Heat induced antigen retrieval with a pH 6.0 sodium citrate buffer is recommended. Optimal concentrations and conditions may vary. |
| <b>Neutralization</b>       | To yield one-half maximal inhibition [ND <sub>50</sub> ] of the biological activity of Human Betacellulin (0.1 ng/ml), a concentration of ≤ 0.1 µg/ml of this antibody is required.                                                                                                                                                                                                                                 |
| <b>Formulation</b>          | A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.                                                                                                                                                                                                                                                                                                                                              |
| <b>Reconstitution</b>       | Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.                                                                                                                                                                                                                                                                                                                |
| <b>Storage</b>              | -20°C                                                                                                                                                                                                                                                                                                                                                                                                               |
| <b>Precautions</b>          | Anti-Human Betacellulin Antibody is for research use only and not for use in diagnostic or therapeutic procedures.                                                                                                                                                                                                                                                                                                  |

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.