

# Anti-Human CXCL16 Antibody

Catalog # ABG10057

## Product Information

---

<b>Application</b>	WB, IHC, E
<b>Reactivity</b>	Human
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal

## Additional Information

---

<b>Preparation</b>	Produced from sera of rabbits pre-immunized with highly pure (>98%) recombinant hCXCL16. Anti-Human CXCL16 specific antibody was purified by affinity chromatography employing immobilized hCXCL16 matrix.
<b>WesternBlot</b>	To detect hCXCL16 by Western Blot analysis this antibody can be used at a concentration of 0.1- 0.2 µg/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant hCXCL16 is 1.5-3.0 ng/lane, under either reducing or non-reducing conditions.
<b>Sandwich</b>	To detect hCXCL16 by sandwich ELISA (using 100 µl/well antibody solution) a concentration of 0.5 - 2.0 µg/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with BioGems' Biotinylated Anti-Human CXCL16 (60-109BT) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant hCXCL16.
<b>Immunohistochemistry</b>	<p>This antibody stained formalin-fixed paraffin-embedded sections of human colorectal adenocarcinoma tissue. The recommended concentration is 0.5 µg/ml with an overnight incubation at 4°C. An HRP-labeled polymer detection system was used with a DAB chromogen. Antigen retrieval is not required for these conditions. Optimal concentrations and conditions may vary.</p> <p>*Additional Immunostaining data available. Please contact Tech Support for information.</p>
<b>Formulation</b>	A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.
<b>Reconstitution</b>	Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.
<b>Storage</b>	-20°C
<b>Precautions</b>	Anti-Human CXCL16 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.