

Anti-Human Eotaxin-2 Antibody

Catalog # ABG10091

Product Information

Application	WB, IHC, E
Reactivity	Human
Host	Mouse
Clonality	Monoclonal

Additional Information

Preparation	Produced in BALB/c mice using highly pure (>98%) recombinant human Eotaxin-2 as the immunizing antigen. This IgG1 _K antibody was purified from ascites fluid by antigen affinity chromatography.
WesternBlot	To detect hEotaxin-2 by Western Blot analysis this antibody can be used at a concentration of 0.25-0.50 μ g/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant hEotaxin-2 is 0.5-1.0 ng/lane, under non-reducing conditions.
Sandwich	In a sandwich ELISA (assuming 100 μ l/well), a concentration of 2.0-4.0 μ g/ml of this antibody will detect at least 80.0 pg/well of recombinant human Eotaxin-2 when used with BioGems' biotinylated antigen affinity purified anti-human Eotaxin-2 (60-123BT) as the detection antibody at a concentration of approximately 1.0-2.0 μ g/ml.
Immunohistochemistry	This antibody stained formalin-fixed, paraffin-embedded sections of human tonsil. The recommended concentration is 10.0 μ g/ml with an overnight incubation at 4 °C. An HRP-labeled polymer detection system was used with a DAB chromogen. Heat induced antigen retrieval with a pH 8.0 EDTA buffer is recommended. Optimal concentrations and conditions may vary. Protocol and staining provided by Dr. Lauren Binge, Laboratory of Prof. Charles Mackay, Monash Univeristy, Australia.
Formulation	A sterile filtered antibody solution was lyophilized from PBS.
Reconstitution	Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.
Storage	-20°C
Precautions	Anti-Human Eotaxin-2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.