

Anti-Human Eotaxin-3 Antibody

Catalog # ABG10094

Product Information

Application	WB, E
Reactivity	Human
Host	Mouse
Clonality	Monoclonal

Additional Information

Preparation	Produced in BALB/c ICRR F ₁ mice using highly pure (>98%) recombinant human Eotaxin-3 as the immunizing antigen. This IgG2 _{ak} antibody was purified from ascites fluid by Protein A affinity chromatography.
WesternBlot	To detect hEotaxin-3 by Western Blot analysis this antibody can be used at a concentration of 0.25-0.50 μ g/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant hEotaxin-3 is 0.25-0.5 ng/lane, under reducing or non-reducing conditions.
Sandwich	In a sandwich ELISA (assuming 100 μ /well), a concentration of 1.0-2.0 μ g/ml of this antibody will detect at least 1000 pg/ml of recombinant human Eotaxin-3 when used with BioGems's biotinylated antigen affinity purified anti-human Eotaxin-3 (60-124BT) as the detection antibody at a concentration of approximately 0.25-0.50 μ g/ml.
Neutralization	To yield one-half maximal inhibition [ND ₅₀] of the biological activity of Human Eotaxin-3 (2.0 μ g/ml), a concentration of 1.0-5.0 μ g/ml of this antibody is required.
Formulation	A sterile filtered antibody solution was lyophilized from PBS.
Reconstitution	Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.
Storage	-20°C
Precautions	Anti-Human Eotaxin-3 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.