

Anti-Murine FGF-9 Antibody

Catalog # ABG10110

Product Information

Application	WB, E
Reactivity	Mouse
Host	Rabbit
Clonality	Polyclonal

Additional Information

Preparation	Produced from sera of rabbits pre-immunized with highly pure (>98%) recombinant mFGF-9. Anti-Murine FGF-9 specific antibody was purified by affinity chromatography employing immobilized mFGF-9 matrix.
WesternBlot	To detect mFGF-9 by Western Blot analysis this antibody can be used at a concentration of 0.1-0.2 μ g/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant mFGF-9 is 1.5-3.0 ng/lane, under either reducing or non-reducing conditions.
Sandwich	To detect mFGF-9 by sandwich ELISA (using 100 μ l/well antibody solution) a concentration of 0.5 - 2.0 μ g/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with BioGems's Biotinylated Anti-Murine FGF-9 (61-069BT) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant mFGF-9.
Neutralization	To yield one-half maximal inhibition [ND50] of the biological activity of mFGF-9 (1.50 ng/ml), a concentration of 0.025-0.06 μ g/ml of this antibody is required.
Formulation	A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.
Reconstitution	Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.
Storage	-20°C
Precautions	Anti-Murine FGF-9 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.