

Anti-Human IGF-II Antibody

Catalog # ABG10186

Product Information

Application	WB, IHC, E
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal

Additional Information

Preparation	Produced from sera of rabbits pre-immunized with highly pure (>98%) recombinant hIGF-II (human IGF-II). Anti-Human IGF-II specific antibody was purified by affinity chromatography employing immobilized hhIGF-II matrix.
WesternBlot	To detect hIGF-II by Western Blot analysis this antibody can be used at a concentration of 0.1- 0.2 μ g/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant hIGF-II is 1.5-3.0 ng/lane, under either reducing or non-reducing conditions.
Sandwich	To detect hIGF-II by sandwich ELISA (using 100 μ l/well antibody solution) a concentration of 0.5 - 2.0 μ g/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with BioGems's Biotinylated Anti-Human IGF-II (60-187BT) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant hIGF-II.
Immunohistochemistry	This antibody stained formalin-fixed, paraffin-embedded sections of normal human placenta. The recommended concentration ranges from 0.5 μ g/ml-5.0 μ g/ml. An HRP-labeled polymer detection system was used with DAB chromogen. Recommended antigen retrievals are heat induced antigen retrieval with a pH 6.0 Sodium Citrate buffer or enzyme induced antigen retrieval with Proteinase K. Optimal concentrations and conditions may vary.
Neutralization	To yield one-half maximal inhibition [ND_{50}] of the biological activity of hIGF-II (10 ng/ml), a concentration of 3.1 - 4.6 μ g/ml of this antibody is required.
Formulation	A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.
Reconstitution	Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.
Storage	-20°C
Precautions	Anti-Human IGF-II Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.