

Anti-Murine IL-2 Antibody

Catalog # ABG10256

Product Information

Application	WB, E
Reactivity	Mouse
Host	Rat
Clonality	Monoclonal

Additional Information

Preparation	Produced in Sprague Dawley rats using highly pure (>98%) recombinant murine IL-2 as the immunizing antigen. This IgG2A _k antibody was purified from cell culture by Protein A affinity chromatography.
WesternBlot	To detect mIL-2 by Western Blot analysis this antibody can be used at a concentration of 2.0-4.0 μ g/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant mIL-2 is 1.0-2.0 ng/lane, under reducing conditions and 0.2-0.4 ng/lane, under non-reducing conditions.
Sandwich	In a sandwich ELISA (assuming 100 μ /well), a concentration of 4.0-8.0 μ g/ml of this antibody will detect at least 32 pg/ml of recombinant murine IL-2 when used with BioGems's biotinylated antigen affinity purified anti-murine IL-2 (61-002BT) as the detection antibody at a concentration of at least 1.0-2.0 μ g/ml.
Formulation	A sterile filtered antibody solution was lyophilized from PBS.
Reconstitution	Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.
Storage	-20°C
Precautions	Anti-Murine IL-2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.