

Anti-Human IL-6 Antibody

Catalog # ABG10290

Product Information

Application	WB, IHC, E
Reactivity	Human
Host	Goat
Clonality	Polyclonal

Additional Information

Preparation	Produced from sera of goats pre-immunized with highly pure recombinant Human IL-6. Anti-Human IL-6 specific antibody was purified by affinity chromatography employing immobilized Human IL-6 matrix.
WesternBlot	To detect Human IL-6 by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 μ g/ml. When used in conjunction with compatible secondary reagents the detection limit for recombinant Human IL-6 is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.
Sandwich	To detect Human IL-6 by sandwich ELISA (using 100 μ l/well antibody solution) a concentration of 0.5 - 2.0 μ g/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with BioGems's Biotinylated Anti-Human IL-6 (60-006GBT) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant Human IL-6.
Immunohistochemistry	This antibody stained formalin-fixed, paraffin-embedded sections of human cervical squamous cell carcinoma. The recommended concentrations are 0.25 μ g/ml with an overnight incubation at 4°C. An HRP-labeled polymer detection system was used with a DAB chromogen. The recommended antigen retrieval is heat induced antigen retrieval with a pH 6.0 Sodium Citrate buffer. Optimal concentrations and conditions may vary.
Neutralization	To yield one-half maximal inhibition [ND ₅₀] of the biological activity of Human IL-6 (5 ng/ml), a concentration of 0.05 - 0.08 μ g/ml of this antibody is required.
Formulation	A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.
Reconstitution	Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.
Storage	-20°C
Precautions	Anti-Human IL-6 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.