

Anti-Rat IL-6 Antibody

Catalog # ABG10293

Product Information

Application	WB, IHC, E
Reactivity	Rat
Host	Rabbit
Clonality	Polyclonal

Additional Information

Preparation	Produced from sera of rabbits pre-immunized with highly pure recombinant Rat IL-6. Anti-Rat IL-6 specific antibody was purified by affinity chromatography employing immobilized Rat IL-6 matrix.
WesternBlot	To detect Rat IL-6 by Western Blot analysis this antibody can be used at a concentration of 0.1- 0.2 μ g/ml. When used in conjunction with compatible secondary reagents, the detection limit for recombinant Rat IL-6 is 1.5-3.0 ng/lane, under either reducing or non-reducing conditions.
Sandwich	To detect Rat IL-6 by sandwich ELISA (using 100 μ l/well antibody solution) a concentration of 0.5 - 2.0 μ g/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with BioGems' Biotinylated Anti-Rat IL-6 (62-006BT) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant Rat IL-6.
Immunohistochemistry	This antibody stained colchicine injected rat brain (cingulate cortex) tissue. The primary antibody was incubated at 1.0 μ g/ml overnight at 4 °C. This was followed by a peroxidase conjugated secondary antibody and then a fluorescein Tyramide Signal Amplification (TSA™) reagent. Optimal concentrations and conditions may vary. Information and photo are courtesy of the Tissue Profiling group, SciLifeLab Stockholm.
Neutralization	To yield one-half maximal inhibition [ND ₅₀] of the biological activity of Rat IL-6 (0.5 ng/ml), a concentration of 1.0-1.50 μ g/ml of this antibody is required.
Formulation	A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.
Reconstitution	Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.
Storage	-20°C
Precautions	Anti-Rat IL-6 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.