

## Anti-Human IL-6 Antibody

Catalog # ABG10300

## **Product Information**

ApplicationWB, IHC, EReactivityHumanHostMouseClonalityMonoclonal

## **Additional Information**

**Preparation** Produced in BALB/c mice using highly pure (>98%) recombinant human IL-6

as the immunizing antigen. This IgG1<sub>K</sub> antibody was purified from cell culture

by Protein A affinity chromatography.

**WesternBlot** To detect hIL-6 by Western Blot analysis this antibody can be used at a

concentration of 0.20-0.40 [g/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant hIL-6 is 0.5-1.0

ng/lane, under reducing or non-reducing conditions.

Sandwich In a sandwich ELISA (assuming 100 □/well), a concentration of 4.0-8.0 □g/ml

of this antibody will detect at least 500 pg/ml of recombinant human IL-6 when used with BioGems' biotinylated antigen affinity purified anti-human IL-6 (60-006BT) as the detection antibody at a concentration of

approximately 0.5-1.0 ☐g/ml.

**Immunohistochemistry**This antibody stained formalin-fixed, paraffin-embedded sections of human

cervical squamous cell carcinoma. The recommended concentration is 0.125  $\mu g/ml$  with an overnight incubation at 4°C. An HRP-labeled polymer detection system was used with a DAB chromogen. Heat induced antigen retrieval with a pH 6.0 sodium citrate buffer is recommended. Optimal concentrations and conditions may vary. Tissue samples were provided by the Cooperative Human Tissue Network, which is funded by the National Cancer Institute.

\*Additional Immunostaining data available. Please contact Tech Support for

information.

**Formulation** A sterile filtered antibody solution was lyophilized from PBS.

**Reconstitution** Centrifuge vial prior to opening. Reconstitute in sterile water to a

concentration of 0.1-1.0 mg/ml.

Storage -20°C

**Precautions** Anti-Human IL-6 Antibody is for research use only and not for use in

diagnostic or therapeutic procedures.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.