

Anti-Rat IP-10 Antibody

Catalog # ABG10317

Product Information

ApplicationWB, IHC, EReactivityRatHostRabbitClonalityPolyclonal

Additional Information

Preparation Produced from sera of rabbits immunized with highly pure recombinant Rat

IP-10. Anti-Rat IP-10 specific antibody was purified by affinity chromatography

employing an immobilized Rat IP-10 matrix.

WesternBlot To detect Rat IP-10 by Western Blot analysis this antibody can be used at a

concentration of 0.1 - 0.2 μ g/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant Rat IP-10 is 1.5 - 3.0

ng/lane, under either reducing or non-reducing conditions.

Sandwich To detect Rat IP-10 by sandwich ELISA (using 100µl/well antibody solution) a

concentration of $0.5 - 2.0 \,\mu\text{g/ml}$ of this antibody is required. This antigen affinity purified antibody, in conjunction with BioGems' Biotinylated Anti-Rat IP-10 (62-062BT) as a detection antibody, allows the detection of at

least 0.2 - 0.4 ng/well of recombinant Rat IP-10.

ImmunohistochemistryThis antibody stained colchicine injected rat brain (including the caudate

putamen) tissue. The primary antibody was incubated at 0.25 μg/ml overnight at 4°C. This was followed by a peroxidase conjugated secondary antibody and then a fluorescein Tyramide Signal Amplification (TSA[™])

reagent. Optimal concentrations and conditions may vary.

Neutralization To yield one-half maximal inhibition [ND_x] of the biological activity of Rat IP-10

(100 ng/ml), a concentration of 5.0-10.0 µg/ml of this antibody is required.

Formulation A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.

Reconstitution Centrifuge vial prior to opening. Reconstitute in sterile water to a

concentration of 0.1-1.0 mg/ml.

Storage -20°C

PrecautionsAnti-Rat IP-10 Antibody is for research use only and not for use in diagnostic

or therapeutic procedures.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.