

Anti-Human Maspin Antibody

Catalog # ABG10350

Product Information

Application	WB, IHC, E
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal

Additional Information

Preparation	Produced from sera of rabbits immunized with highly pure recombinant Human Maspin. Anti-Human Maspin specific antibody was purified by affinity chromatography employing an immobilized Human Maspin matrix.
WesternBlot	To detect hMaspin by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 µg/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant hMaspin is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.
Sandwich	To detect hMaspin by sandwich ELISA (using 100µl/well antibody solution) a concentration of 0.5 - 2.0 µg/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with BioGems's Biotinylated Anti-Human Maspin (60-210BT) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant hMaspin.
Immunohistochemistry	This antibody stained formalin-fixed paraffin-embedded sections of human normal breast and human breast lobular carcinoma in situ. The recommended concentration is 7.8 ng/mL- 15.625 ng/mL with an overnight incubation at 4°C. An HRP-labeled polymer detection system was used with a DAB chromogen. Optimal results for these conditions were achieved without antigen retrieval. Optimal concentrations and conditions may vary.
Formulation	A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.
Reconstitution	Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.
Storage	-20°C
Precautions	Anti-Human Maspin Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.