

Anti-Human MCP-2 Antibody

Catalog # ABG10357

Product Information

Application	WB, IHC, E
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal

Additional Information

Preparation	Produced from sera of rabbits pre-immunized with highly pure (>98%) recombinant hMCP-2. Anti-Human MCP-2 specific antibody was purified by affinity chromatography employing immobilized hMCP-2 matrix.
WesternBlot	To detect hMCP-2 by Western Blot analysis this antibody can be used at a concentration of 0.1-0.2 µg/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant hMCP-2 is 1.5-3.0 ng/lane, under either reducing or non-reducing conditions.
Sandwich	To detect hMCP-2 by sandwich ELISA (using 100 µl/well antibody solution) a concentration of 0.5 - 2.0 µg/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with BioGems' Biotinylated Anti-Human MCP-2 (60-213BT) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant hMCP-2.
Immunohistochemistry	This antibody stained formalin-fixed, paraffin-embedded sections of human normal gastric mucosa. The recommended concentration is 0.75mg/ml with an overnight incubation at 4 °C. An HRP-labeled polymer detection system was used with a DAB chromogen. Heat induced antigen retrieval with a pH 6.0 sodium citrate buffer is recommended. Optimal concentrations and conditions may vary.
Tissue samples were provided by the Cooperative Human Tissue Network, which is funded by the National Cancer Institute.	
Formulation	A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.
Reconstitution	Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.
Storage	-20°C
Precautions	Anti-Human MCP-2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.