

# Biotinylated Anti-Human NT-4 Antibody

Catalog # ABG10447

## Product Information

---

<b>Application</b>	WB, E
<b>Reactivity</b>	Human
<b>Host</b>	Goat
<b>Clonality</b>	Polyclonal

## Additional Information

---

<b>Preparation</b>	Produced from sera of goats pre-immunized with highly pure (>98%) recombinant hNT-4. Anti-Human NT-4 specific antibody was purified by affinity chromatography and then biotinylated.
<b>WesternBlot</b>	To detect hNT-4 by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 µg/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant hNT-4 is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.
<b>Sandwich</b>	To detect hNT-4 by sandwich ELISA (using 100 µl/well antibody solution) a concentration of 0.25 – 1.0 µg/ml of this antibody is required. This biotinylated polyclonal antibody, in conjunction with BioGems's Polyclonal Anti-Human NT-4 (60-253P) as a capture antibody, allows the detection of at least 0.2 – 0.4 ng/well of recombinant hNT-4.
<b>Direct</b>	To detect hNT-4 by direct ELISA (using 100 µl/well antibody solution) a concentration of 0.25 – 1.0 µg/ml of this antibody is required. This biotinylated polyclonal antibody, in conjunction with compatible secondary reagents, allows the detection of at least 0.2 – 0.4 ng/well of recombinant hNT-4.
<b>Formulation</b>	A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.
<b>Reconstitution</b>	Centrifuge vial prior to opening. Reconstitute in sterile PBS containing 0.1% BSA to a concentration of 0.1-1.0 mg/ml.
<b>Storage</b>	-20°C
<b>Precautions</b>	Biotinylated Anti-Human NT-4 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.