

Anti-Rat SCF Antibody

Catalog # ABG10492

Product Information

Application	WB, E
Reactivity	Rat
Host	Rabbit
Clonality	Polyclonal

Additional Information

Preparation	Produced from sera of rabbits immunized with highly pure recombinant Rat SCF. Anti-Rat SCF specific antibody was purified by affinity chromatography employing an immobilized Rat SCF matrix.
WesternBlot	To detect rat SCF by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 µg/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant rat SCF is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.
Sandwich	To detect rat SCF by sandwich ELISA (using 100µl/well antibody solution) a concentration of 0.5 - 2.0 µg/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with BioGems's Biotinylated Anti-Rat SCF (62-073BT) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant rat SCF.
Neutralization	To yield one-half maximal inhibition [ND ₅₀] of the biological activity of rat SCF (50.0 ng/ml), a concentration of 1.2-1.8 µg/ml of this antibody is required.
Formulation	A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.
Reconstitution	Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.
Storage	-20°C
Precautions	Anti-Rat SCF Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.