

Anti-Human TACI Antibody

Catalog # ABG10529

Product Information

Application	WB, E
Reactivity	Human
Host	Goat
Clonality	Polyclonal

Additional Information

Preparation	Produced from sera of goats pre-immunized with highly pure (>98%) recombinant h-TACI. Anti-Human TACI specific antibody was purified by affinity chromatography employing immobilized h-TACI matrix.
WesternBlot	To detect hTACI by Western Blot analysis this antibody can be used at a concentration of 0.1- 0.2 µg/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant h-TACI is 1.5-3.0 ng/lane, under either reducing or non-reducing conditions.
Sandwich	To detect hTACI by sandwich ELISA (using 100 µl/well antibody solution) a concentration of 0.5 - 2.0 µg/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with BioGems' Biotinylated Anti-Human TACI (60-302BT) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant hTACI.
Formulation	A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.
Reconstitution	Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.
Storage	-20°C
Precautions	Anti-Human TACI Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.