

## Anti-Human TECK Antibody

Catalog # ABG10535

## **Product Information**

ApplicationWB, IHC, EReactivityHumanHostMouseClonalityMonoclonal

## **Additional Information**

**Preparation** Produced in BALB/c x ICR F<sub>1</sub> mice using highly pure (>98%) recombinant

human TECK as the immunizing antigen. This IgG1<sub>K</sub> antibody was purified

from ascites fluid by Protein A affinity chromatography.

WesternBlot To detect hTECK by Western Blot analysis this antibody can be used at a

concentration of 0.25-0.50 [g/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant hTECK is 0.50-1.0

ng/lane, under non-reducing conditions.

Sandwich In a sandwich ELISA (assuming 100 □/well), a concentration of 1.0-2.0 □g/ml

of this antibody will detect at least 300 pg/ml of recombinant human TECK when used with BioGems' biotinylated antigen affinity purified anti-human TECK (60-305BT) as the detection antibody at a concentration of

approximately 0.5-1.0 ☐g/ml.

**Immunohistochemistry**This antibody stained formalin-fixed, paraffin-embedded sections of human

normal small bowel. The recommended concentration is 2.5 [g/mL with an overnight incubation at 4°C. An HRP-labeled polymer detection system was used with a DAB chromogen and a Proteinase K antigen retrieval. Optimal

concentrations and conditions may vary.

Tissue samples were provided by the Cooperative Human Tissue Network, which is funded by the National Cancer Institute. Information and photo are courtesy of Flagship Biosciences, LLC – Histology Services.

**Formulation** A sterile filtered antibody solution was lyophilized from PBS.

**Reconstitution** Centrifuge vial prior to opening. Reconstitute in sterile water to a

concentration of 0.1-1.0 mg/ml.

Storage -20°C

**Precautions**Anti-Human TECK Antibody is for research use only and not for use in

diagnostic or therapeutic procedures.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.