

# Anti-Murine TNF- $\alpha$ Antibody

Catalog # ABG10547

## Product Information

---

|                    |            |
|--------------------|------------|
| <b>Application</b> | WB, IHC, E |
| <b>Reactivity</b>  | Mouse      |
| <b>Host</b>        | Rabbit     |
| <b>Clonality</b>   | Polyclonal |

## Additional Information

---

|                             |                                                                                                                                                                                                                                                                                                                                                                                                                                                               |
|-----------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Preparation</b>          | Produced from sera of rabbits pre-immunized with highly pure recombinant Murine TNF- $\alpha$ . Anti-Murine TNF- $\alpha$ specific antibody was purified by affinity chromatography employing immobilized Murine TNF- $\alpha$ matrix.                                                                                                                                                                                                                        |
| <b>WesternBlot</b>          | To detect Murine TNF- $\alpha$ by Western Blot analysis this antibody can be used at a concentration of 0.1-0.2 $\mu$ g/ml. When used in conjunction with compatible secondary reagents, the detection limit for recombinant Murine TNF- $\alpha$ is 1.5-3.0 ng/lane, under either reducing or non-reducing conditions.                                                                                                                                       |
| <b>Sandwich</b>             | To detect Murine TNF- $\alpha$ by sandwich ELISA (using 100 $\mu$ l/well antibody solution) a concentration of 0.5 - 2.0 $\mu$ g/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with BioGems' Biotinylated Anti-Murine TNF- $\alpha$ (61-124BT) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant Murine TNF- $\alpha$ .                                                   |
| <b>Immunohistochemistry</b> | This antibody stained colchicine injected mouse brain (hippocampus CA1 region) tissue. The primary antibody was incubated at 1.0 $\mu$ g/ml overnight at 4 °C. This was followed by a peroxidase conjugated secondary antibody and then a fluorescein Tyramide Signal Amplification (TSA™) reagent. Optimal concentrations and conditions may vary.<br><small>Information and photo are courtesy of the Tissue Profiling group, SciLifeLab Stockholm.</small> |
| <b>Neutralization</b>       | To yield one-half maximal inhibition [ND <sub>50</sub> ] of the biological activity of Murine TNF- $\alpha$ (0.5 ng/ml), a concentration of 0.04 - 0.05 $\mu$ g/ml of this antibody is required.                                                                                                                                                                                                                                                              |
| <b>Formulation</b>          | A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.                                                                                                                                                                                                                                                                                                                                                                                        |
| <b>Reconstitution</b>       | Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.                                                                                                                                                                                                                                                                                                                                                          |
| <b>Storage</b>              | -20°C                                                                                                                                                                                                                                                                                                                                                                                                                                                         |
| <b>Precautions</b>          | Anti-Murine TNF- $\alpha$ Antibody is for research use only and not for use in diagnostic or therapeutic procedures.                                                                                                                                                                                                                                                                                                                                          |

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.