10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Anti-Human sTRAIL/Apo2L Antibody

Catalog # ABG10566

Product Information

Application	WB, IHC, E
Reactivity	Human
Host	Mouse
Clonality	Monoclonal

Additional Information

Preparation

Produced in BALB/c x ICR F_1 mice using highly pure (>98%) recombinant human TRAIL/APO2L as the immunizing antigen. This $IgG1_K$ antibody was purified from ascites fluid by Protein A affinity chromatography.

WesternBlot

To detect Human sTRAIL/Apo2L by Western Blot analysis this antibody can be used at a concentration of 0.25-0.50 g/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant Human sTRAIL/Apo2L is 0.25-1.0 ng/lane, under reducing or non-reducing conditions.

Sandwich

In a sandwich ELISA (assuming 100 l/well), a concentration of 2.0-4.0 g/ml of this antibody will detect at least 500 pg/ml of recombinant human sTRAIL/APO2L when used with BioGems' biotinylated antigen affinity purified anti-human sTRAIL/APO2L (60-330BT) as the detection antibody at a concentration of approximately 1.0-2.0 g/ml.

Immunohistochemistry

This antibody stained formalin-fixed, paraffin-embedded sections of human renal cell carcinoma. The recommended concentration is 2.5 g/mL- 1.5 g/mL with an overnight incubation at 4˚C. An HRP-labeled polymer detection system was used with a DAB chromogen. Heat induced antigen retrieval with a pH 9.0 buffer is recommended. Optimal concentrations and conditions may vary.

Tissue samples were provided by the Cooperative Human Tissue Network, which is funded by the National Cancer Institute.
Information and photo are courtesy of Flagship Biosciences, LLC – Histology Services.

Neutralization

To yield one-half maximal inhibition [ND_{50}] of the biological activity of Human TRAIL/Apo2L (10.0 ng/ml), a concentration of 0.30-0.60 µg/ml of this antibody is required.

Formulation

A sterile filtered antibody solution was lyophilized from PBS.

Reconstitution

Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.

Storage

-20°C

Precautions

Anti-Human sTRAIL/Apo2L Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.