

Anti-CD11c Picoband Antibody

Catalog # ABO10065

Product Information

Application	WB, IHC-P, E
Primary Accession	P20702
Host	Rabbit
Reactivity	Human, Mouse
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for CD11c detection. Tested with WB, IHC-P, Direct ELISA in Human;Mouse.

Reconstitution Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	3687
Other Names	Integrin alpha-X, CD11 antigen-like family member C, Leu M5, Leukocyte adhesion glycoprotein p150, 95 alpha chain, Leukocyte adhesion receptor p150, 95, CD11c, ITGAX, CD11C
Calculated MW	127829
Application Details	Western blot, 0.1-0.5 μ g/ml Immunohistochemistry(Paraffin-embedded Section), 0.5-1 μ g/ml Direct ELISA, 0.1-0.5 μ g/ml
Subcellular Localization	Membrane; Single-pass type I membrane protein.
Tissue Specificity	Predominantly expressed in monocytes and granulocytes.
Source	Eukaryota
Contents	Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg NaN ₃ .
Immunogen	E. coli-derived human CD11c recombinant protein (Position: S161-T342).
Cross Reactivity	No cross reactivity with other proteins.
Storage	At -20 °C; for one year. After r ° Constitution, at 4 °C; for one month. It ° Can also be aliquotted and stored frozen at -20 °C; for a longer time. Avoid repeated freezing and thawing.

Protein Information

Name	ITGAX (HGNC:6152)
Synonyms	CD11C
Function	Integrin alpha-X/beta-2 is a receptor for fibrinogen. It recognizes the sequence G-P-R in fibrinogen. It mediates cell-cell interaction during inflammatory responses. It is especially important in monocyte adhesion and chemotaxis. Functions as a receptor of the erythrocyte-specific adhesion molecule ICAM4 and mediates erythrophagocytosis (PubMed: 16985175).
Cellular Location	Cell membrane; Single-pass type I membrane protein
Tissue Location	Predominantly expressed in monocytes and granulocytes

Background

CD11c, also known as Integrin, alpha X (complement component 3 receptor 4 subunit) (ITGAX), is a gene that encodes for CD11c. This gene encodes the integrin alpha X chain protein. Integrins are heterodimeric integral membrane proteins composed of an alpha chain and a beta chain. This protein combines with the beta 2 chain (ITGB2) to form a leukocyte-specific integrin referred to as inactivated-C3b (iC3b) receptor 4 (CR4). The alpha X beta 2 complex seems to overlap the properties of the alpha M beta 2 integrin in the adherence of neutrophils and monocytes to stimulated endothelium cells, and in the phagocytosis of complement coated particles. Two transcript variants encoding different isoforms have been found for this gene.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.