

Anti-Cytokeratin 5 Picoband Antibody

Catalog # ABO10071

Product Information

Application	WB, IHC-P
Primary Accession	P13647
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for Keratin, type II cytoskeletal 5(KRT5) detection. Tested with WB, IHC-P in Human;Mouse;Rat.
Reconstitution	Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	3852
Other Names	Keratin, type II cytoskeletal 5, 58 kDa cytokeratin, Cytokeratin-5, CK-5, Keratin-5, K5, Type-II keratin Kb5, KRT5
Calculated MW	62378
Application Details	Immunohistochemistry(Paraffin-embedded Section), 0.5-1 µg/ml, Human, Mouse, Rat, By Heat Western blot, 0.1-0.5 µg/ml, Human, Mouse
Source	Eukaryota
Protein Name	Keratin, type II cytoskeletal 5
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg NaN ₃ .
Immunogen	A synthetic peptide corresponding to a sequence in the middle region of human Cytokeratin 5 (286-317aa KVELEAKVDALMDEINFMKMFFDAELSQQMTH), different from the related mouse sequence by one amino acid, and identical to the related rat sequence.
Purification	Immunogen affinity purified.
Cross Reactivity	No cross reactivity with other proteins
Storage	At -20 °C for one year. After reconstitution, at 4 °C for one month. It can also be aliquotted and stored frozen at -20 °C for a longer time. Avoid repeated freezing and thawing.


Protein Information

Name	KRT5 (HGNC:6442)
Function	Structural component of intermediate filaments in basal keratinocytes of stratified epithelia. Together with its obligate type I partner KRT14, contributes to the formation of the keratin intermediate filament network that provides mechanical stability and resilience to the basal layer of the epidermis (PubMed: 1372711 , PubMed: 1694855). Regulates the recruitment of Langerhans cells to the epidermis, potentially by modulation of the abundance of macrophage chemotactic cytokines, macrophage inflammatory cytokines and CTNND1 localization in keratinocytes (By similarity).
Cellular Location	Cytoplasm.
Tissue Location	Expressed in corneal epithelium (at protein level) (PubMed:26758872). Expressed in keratinocytes (at protein level) (PubMed:20128788, PubMed:31302245).


Background

Cytokeratin 5, also known as KRT5, K5, or CK5, is a protein that is encoded in humans by the KRT5 gene. The protein encoded by this gene is a member of the keratin gene family. The type II cytokeratins consist of basic or neutral proteins which are arranged in pairs of heterotypic keratin chains coexpressed during differentiation of simple and stratified epithelial tissues. This type II cytokeratin is specifically expressed in the basal layer of the epidermis with family member KRT14. Mutations in these genes have been associated with a complex of diseases termed epidermolysis bullosa simplex. The type II cytokeratins are clustered in a region of chromosome 12q12-q13.


Images



Western blot analysis of Cytokeratin 5 expression in rat kidney extract (lane 1), COLO320 whole cell lysates (lane 2) and HELA whole cell lysates (lane 3). Cytokeratin 5 at 62KD was detected using rabbit anti- Cytokeratin 5 Antigen Affinity purified polyclonal antibody (Catalog # ABO10071) at 0.5 μ g/mL. The blot was developed using chemiluminescence (ECL) method .



Cytokeratin 5 was detected in paraffin-embedded sections of human tonsil tissues using rabbit anti- Cytokeratin 5 Antigen Affinity purified polyclonal antibody (Catalog # ABO10071) at 1 μ g/mL. The immunohistochemical section was developed using SABC method .



Cytokeratin 5 was detected in paraffin-embedded sections of human oesophagus squama cancer tissues using rabbit anti- Cytokeratin 5 Antigen Affinity purified polyclonal antibody (Catalog # ABO10071) at 1 μ g/mL. The immunohistochemical section was developed using SABC method .

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.