

Anti-splicing factor 1 Picoband Antibody

Catalog # ABO10133

Product Information

Application	WB, IHC-P, E
Primary Accession	A01009
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for splicing factor 1 detection. Tested with WB, IHC-P, Direct ELISA in Human;Mouse;Rat.
Reconstitution	Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Application Details	Western blot, 0.1-0.5 μ g/ml Immunohistochemistry(Paraffin-embedded Section), 0.5-1 μ g/ml Direct ELISA, 0.1-0.5 μ g/ml
Subcellular Localization	Nucleus.
Tissue Specificity	Detected in lung, ovary, adrenal gland, colon, kidney, muscle, pancreas, thyroid, placenta, brain, liver and heart.
Contents	Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg NaN ₃ .
Immunogen	E. coli-derived human splicing factor 1 recombinant protein (Position: R160-Q266).
Cross Reactivity	No cross reactivity with other proteins.
Storage	At -20° C; for one year. After r° Constitution, at 4° C; for one month. It° Can also be aliquotted and stored frozen at -20° C; for a longer time. Avoid repeated freezing and thawing.

Protein Information

Background

Splicing factor 1 also known as zinc finger protein 162 (ZFM162) is a protein that in humans is encoded by the SF1 gene. This gene encodes a nuclear pre-mRNA splicing factor. The encoded protein specifically recognizes the intron branch point sequence at the 3' splice site, together with the large subunit of U2 auxiliary factor (U2AF), and is required for the early stages of spliceosome assembly. It also plays a role in

nuclear pre-mRNA retention and transcriptional repression. The encoded protein contains an N-terminal U2AF ligand motif, a central hnRNP K homology motif and quaking 2 region which bind a key branch-site adenosine within the branch point sequence, a zinc knuckles domain, and a C-terminal proline-rich domain. Alternative splicing results in multiple transcript variants.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.