

Anti-Cytoglobin Picoband Antibody

Catalog # ABO10162

Product Information

Application	WB, IHC-P
Primary Accession	Q8WWM9
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for Cytoglobin(CYGB) detection. Tested with WB, IHC-P in Human;Mouse;Rat.
Reconstitution	Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	114757
Other Names	Cytoglobin, Histoglobin, HGb, Stellate cell activation-associated protein, CYGB, STAP
Calculated MW	21405
Application Details	Immunohistochemistry(Paraffin-embedded Section), 0.5-1 μ g/ml, Human, Mouse, Rat, By Heat Western blot, 0.1-0.5 μ g/ml, Mouse, Rat, Human
Subcellular Localization	Cytoplasm .
Tissue Specificity	Ubiquitously expressed. Highest expression in heart, stomach, bladder and small intestine. .
Source	Eukaryota
Protein Name	Cytoglobin
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg Na ₃ N.
Immunogen	E.coli-derived human Cytoglobin recombinant protein (Position: M1-P190). Human Cytoglobin shares 95.3% and 93.7% amino acid (aa) sequence identity with mouse and rat Cytoglobin, respectively.
Purification	Immunogen affinity purified.
Cross Reactivity	No cross reactivity with other proteins

Storage	At -20 °C for one year. After r° Constitution, at 4 °C for one month. It° Can also be aliquotted and stored frozen at -20 °C for a longer time.Avoid repeated freezing and thawing.
----------------	---

Protein Information


Name	CYGB (HGNC:16505)
Function	Probable multifunctional globin with a hexacoordinated heme iron required for the catalysis of various reactions depending on redox condition of the cell as well as oxygen availability (PubMed: 11893755 , PubMed: 12359339 , PubMed: 15165856 , PubMed: 19147491 , PubMed: 20511233 , PubMed: 28393874 , PubMed: 28671819 , PubMed: 29128400 , PubMed: 33576020 , PubMed: 34930834). Has a nitric oxide dioxygenase (NOD) activity and is most probably involved in cell-mediated and oxygen-dependent nitric oxide consumption (PubMed: 19147491 , PubMed: 20511233 , PubMed: 28393874 , PubMed: 28671819). By scavenging this second messenger may regulate several biological processes including endothelium-mediated vasodilation and vascular tone (PubMed: 19147491 , PubMed: 28393874). Under normoxic conditions functions as a nitric oxide dioxygenase (NOD) but under hypoxic conditions the globin may switch its function to that of a nitrite (NO ₂) reductase (NiR), generating nitric oxide (PubMed: 29128400). Could also have peroxidase and superoxide dismutase activities, detoxifying reactive oxygen species and protecting cells against oxidative stress (PubMed: 12359339 , PubMed: 33576020 , PubMed: 34930834). Also binds dioxygen with low affinity and could function as an oxygen sensor but has probably no function as a respiratory oxygen carrier (PubMed: 11893755 , PubMed: 15299006 , PubMed: 20553503).
Cellular Location	Cytoplasm. Nucleus
Tissue Location	Widely expressed. Highest expression in heart, stomach, bladder and small intestine.

Background

Cytoglobin(CYGB), also called HGB or STAP, is a ubiquitously expressed hexacoordinate hemoglobin that may facilitate diffusion of oxygen through tissues, scavenge nitric oxide or other reactive oxygen species, or serve a protective function during oxidative stress. The cytoglobin gene is mapped on 17q25.1. The CYGB gene contains 4 exons and spans about 9 kb. Cytoglobin has many elements common to vertebrate globins, including invariant histidine residues, and the amino acids that form the heme pocket share similarity with pentacoordinate myoglobin. In contrast to the high oxygen affinities displayed by most hexacoordinate hemoglobins, the characteristics of CYGB indicate that it can facilitate oxygen transport. Because the oxygen affinity of CYGB is more similar to myoglobin than to neuroglobin, and the oxy form of CYGB resists autooxidation, CYGB is proposed to represent a tissue oxygen reservoir.

Images

Figure 1. Western blot analysis of Cytoglobin using anti-Cytoglobin antibody (ABO10162). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions. Lane 1: rat small intestine tissue lysates, Lane 2: mouse small intestine tissue lysates, After Electrophoresis, proteins were transferred to a



Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-Cytochrome c antigen affinity purified polyclonal antibody (Catalog # ABO10162) at 0.5 μ g/mL overnight at 4 $^{\circ}$ C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for Cytochrome c at approximately 21KD. The expected band size for Cytochrome c is at 21KD.




Figure 2. IHC analysis of Cytochrome c using anti- Cytochrome c antibody (ABO10162).Cytochrome c was detected in paraffin-embedded section of mouse intestine tissues. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1 μ g/ml rabbit anti- Cytochrome c Antibody (ABO10162) overnight at 4 $^{\circ}$ C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 $^{\circ}$ C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.




Figure 3. IHC analysis of Cytochrome c using anti- Cytochrome c antibody (ABO10162).Cytochrome c was detected in paraffin-embedded section of mouse cardiac muscle tissues. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1 μ g/ml rabbit anti- Cytochrome c Antibody (ABO10162) overnight at 4 $^{\circ}$ C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 $^{\circ}$ C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.





Figure 4. IHC analysis of Cytochrome c using anti- Cytochrome c antibody (ABO10162).Cytochrome c was detected in paraffin-embedded section of rat intestine tissues. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1 μ g/ml rabbit anti- Cytochrome c Antibody (ABO10162) overnight at 4 $^{\circ}$ C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 $^{\circ}$ C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.

Figure 5. IHC analysis of Cytochrome c using anti- Cytochrome c antibody (ABO10162).Cytochrome c was detected in paraffin-embedded section of human intestinal cancer



tissues. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1 μ g/ml rabbit anti- Cytoglobin Antibody (ABO10162) overnight at 4 $^{\circ}$ C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 $^{\circ}$ C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.




Figure 6. IHC analysis of Cytoglobin using anti- Cytoglobin antibody (ABO10162).Cytoglobin was detected in paraffin-embedded section of human lung cancer tissues. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1 μ g/ml rabbit anti- Cytoglobin Antibody (ABO10162) overnight at 4 $^{\circ}$ C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 $^{\circ}$ C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.