

Anti-VCAM1 Picoband Antibody

Catalog # ABO10163

Product Information

Application	WB, E
Primary Accession	A01199-1
Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for VCAM1 detection. Tested with WB, ELISA(Cap) in Human.

Reconstitution Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Application Details	Western blot, 0.1-0.5 µg/ml ELISA(Cap), 0.1-0.5 µg/ml
Subcellular Localization	Membrane; Single-pass type I membrane protein.
Tissue Specificity	Expressed on inflamed vascular endothelium, as well as on macrophage-like and dendritic cell types in both normal and inflamed tissue.
Contents	Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg NaN ₃ .
Immunogen	E. coli-derived human VCAM1 recombinant protein (Position: F25-L270).
Cross Reactivity	No cross reactivity with other proteins.
Storage	At -20 °C; for one year. After reconstitution, at 4 °C; for one month. It can also be aliquotted and stored frozen at -20 °C; for a longer time. Avoid repeated freezing and thawing.

Protein Information

Background

CD106(cluster of differentiation 106) also known as vascular cell adhesion molecule 1(VCAM-1), is a protein that in humans is encoded by the VCAM1 gene. In inflammatory conditions and in cardiac allografts undergoing rejection, VCAM1 is upregulated in endothelium of postcapillary venules. Arterial expression of VCAM1 is also found in experimental models of atherosclerosis in the rabbit. This gene is mapped to

chromosome 1 by Southern analysis of somatic cell hybrids. VCAM-1 functions as a cell adhesion molecule. The VCAM-1 protein mediates the adhesion of lymphocytes, monocytes, eosinophils, and basophils to vascular endothelium. It also functions in leukocyte-endothelial cell signal transduction, and it may play a role in the development of atherosclerosis and rheumatoid arthritis. CAM741 works similar to cotransin in that it represses the biosynthesis of VCAM1 cells by blocking the process of cotranslational translocation, which is dependent on the signal peptide of VCAM1. Among the lung metastasis signature genes identified, several, including VCAM1, were functionally validated.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.