

Anti-TRPC4 Picoband Antibody

Catalog # ABO10227

Product Information

Application	WB
Primary Accession	A02037-1
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for TRPC4 detection. Tested with WB in Human;Mouse;Rat.

Reconstitution Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Application Details	Western blot, 0.1-0.5 µg/ml
Subcellular Localization	Membrane; Enhanced insertion into the cell membrane after activation of the EGF receptor.
Tissue Specificity	Strongly expressed in placenta. Expressed at lower levels in heart, pancreas, kidney and brain. Expressed in endothelial cells. Isoform alpha was found to be the predominant isoform. Isoform beta was not found in pancreas and brain.
Contents	Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg NaN ₃ .
Immunogen	A synthetic peptide corresponding to a sequence of human TRPC4 (MAQFYKRNVNAPYRDRIPLRIVRAESEL).
Cross Reactivity	No cross reactivity with other proteins.
Storage	At -20° C; for one year. After reconstitution, at 4° C; for one month. It can also be aliquotted and stored frozen at -20° C; for a longer time. Avoid repeated freezing and thawing.

Protein Information

Background

The short transient receptor potential channel 4 (TRPC4), also known as Trp-related protein 4, is a protein that in humans is encoded by the TRPC4 gene. TRPC4 is a member of the transient receptor potential cation channels. It is mapped to 13q13.3. This protein forms a non-selective calcium-permeable cation channel that

is activated by Gq-coupled receptors and tyrosine kinases, and plays a role in multiple processes including endothelial permeability, vasodilation, neurotransmitter release and cell proliferation. TRPC4 also has an essential role in endothelial-dependent regulation of vascular tone, endothelial permeability, and neurotransmitter release from thalamic interneurons.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.