

Anti-PIAS3 Picoband Antibody

Catalog # ABO10261

Product Information

Application	WB
Primary Accession	Q9Y6X2
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for PIAS3 detection. Tested with WB in Human;Mouse;Rat.

Reconstitution Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	10401
Other Names	E3 SUMO-protein ligase PIAS3, 6.3.2.-, Protein inhibitor of activated STAT protein 3, PIAS3
Calculated MW	68017
Application Details	Western blot, 0.1-0.5 µg/ml
Subcellular Localization	Cytoplasm.
Tissue Specificity	Widely expressed.
Source	Eukaryota
Contents	Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg NaN ₃ .
Immunogen	A synthetic peptide corresponding to a sequence of human PIAS3 (QRFEEAHFTFALTPQQVQQILTSREVLPGAKCDYTIQVQLRF).
Cross Reactivity	No cross reactivity with other proteins.
Storage	At -20 °C; for one year. After r ° Constitution, at 4 °C; for one month. It ° Can also be aliquotted and stored frozen at -20 °C; for a longer time. Avoid repeated freezing and thawing.

Protein Information

Name	PIAS3
-------------	-------

Function	Functions as an E3-type small ubiquitin-like modifier (SUMO) ligase, stabilizing the interaction between UBE2I and the substrate, and as a SUMO-tethering factor. Plays a crucial role as a transcriptional coregulation in various cellular pathways, including the STAT pathway and the steroid hormone signaling pathway. Involved in regulating STAT3 signaling via inhibiting STAT3 DNA-binding and suppressing cell growth. Enhances the sumoylation of MTA1 and may participate in its paralog-selective sumoylation (PubMed: 21965678 , PubMed: 9388184). Sumoylates CCAR2 which promotes its interaction with SIRT1 (PubMed: 25406032). Diminishes the sumoylation of ZFH3 by preventing the colocalization of ZFH3 with SUMO1 in the nucleus (PubMed: 24651376).
Cellular Location	Cytoplasm {ECO:0000250 UniProtKB:O54714}. Nucleus {ECO:0000250 UniProtKB:O54714}. Nucleus speckle {ECO:0000250 UniProtKB:O54714}. Note=Colocalizes with MITF in the nucleus. Colocalizes with GFI1 in nuclear dots. Colocalizes with SUMO1 in nuclear granules. {ECO:0000250 UniProtKB:O54714}
Tissue Location	Widely expressed..

Background

E3 SUMO-protein ligase PIAS3 is an enzyme that in humans is encoded by the PIAS3 gene. This gene encodes a member of the PIAS [protein inhibitor of activated STAT (signal transducer and activator of transcription)] family of transcriptional modulators. The protein functions as a SUMO (small ubiquitin-like modifier)-E3 ligase which catalyzes the covalent attachment of a SUMO protein to specific target substrates. It directly binds to several transcription factors and either blocks or enhances their activity. Alternatively spliced transcript variants of this gene have been identified, but the full-length nature of some of these variants has not been determined.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.