10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

# Anti-CDK1 Antibody (Monoclonal, A17)

Catalog # ABO10462

## Product Information

| | |
|---|---|
| **Application** | WB, IHC-P, IHC-F |
| **Primary Accession** | P39951 |
| **Host** | Mouse |
| **Isotype** | Mouse IgG2a |
| **Reactivity** | Human, Mouse |
| **Clonality** | Monoclonal |
| **Format** | Lyophilized |
| **Description** | Mouse IgG monoclonal antibody for p34cdc2, cyclin-dependent kinase 1 (CDK1) detection. Tested with WB, IHC-P, IHC-F in Human;mouse;chicken. No cross reactivity with other proteins. |
| **Reconstitution** | Add 1ml of PBS buffer will yield a concentration of 100ug/ml. |

## Additional Information

| | |
|---|---|
| **Gene ID** | 54237 |
| **Other Names** | Cyclin-dependent kinase 1, CDK1, 2.7.11.22, 2.7.11.23, Cell division control protein 2 homolog, Cell division protein kinase 1, p34 protein kinase, Cdk1, Cdc2, Cdc2a, Cdkn1 |
| **Calculated MW** | 34135 |
| **Application Details** | Immunohistochemistry(Paraffin-embedded Section), 1-2 □g/ml, Human, chicken, mouse, By Heat<br><br>Immunohistochemistry(Frozen Section), 1-2 □g/ml, Human, chicken, mouse, -<br>Western blot, 0.5-1 □g/ml, Human, chicken, mouse |
| **Subcellular Localization** | Nucleus . Cytoplasm . Mitochondrion . Cytoplasm, cytoskeleton, microtubule organizing center, centrosome . Cytoplasm, cytoskeleton, spindle . Cytoplasmic during the interphase. Reversibly translocated from cytoplasm to nucleus when phosphorylated before G2-M transition when associated with cyclin B1. Accumulates in mitochondria in G2-arrested cells upon DNA-damage. Colocalizes with SIRT2 on centrosome during prophase and on splindle fibers during metaphase of the mitotic cell cycle (By similarity). . |
| **Tissue Specificity** | CDK1: Isoform 2 is found in breast cancer tissues. |
| **Source** | Eukaryota |
| **Protein Name** | Cyclin-dependent kinase 1 |

| | |
|---|---|
| **Contents** | Mouse ascites fluid, 1.2% sodium acetate, 2mg BSA, with 0.01mg NaN3 as preservative. |
| **Clone Names** | A17 |
| **Immunogen** | C-terminal two-thirds of Xenopus CDK1 expressed in E.coli. |
| **Purification** | Ascites |
| **Cross Reactivity** | No cross reactivity with other proteins |
| **Storage** | At -20˚C for one year. After r˚Constitution, at 4˚C for one month. It˚Can also be aliquotted and stored frozen at -20˚C for a longer time.Avoid repeated freezing and thawing. |
| **Sequence Similarities** | Belongs to the protein kinase superfamily. CMGC Ser/Thr protein kinase family. CDC2/CDKX subfamily. |

## Protein Information

| | |
|---|---|
| **Name** | Cdk1 |
| **Synonyms** | Cdc2, Cdc2a, Cdkn1 |
| **Function** | Plays a key role in the control of the eukaryotic cell cycle by modulating the centrosome cycle as well as mitotic onset; promotes G2-M transition via association with multiple interphase cyclins (PubMed:10542199, PubMed:19821535). Phosphorylates PARVA/actopaxin, APC, AMPH, APC, ASB7, BARD1, Bcl-xL/BCL2L1, BRCA2, CALD1, CASP8, CDC7, CDC20, CDC25A, CDC25C, CC2D1A, CENPA, CSNK2 proteins/CKII, FZR1/CDH1, CDK7, CEBPB, CHAMP1, DMD/dystrophin, EEF1 proteins/EF-1, EZH2, KIF11/EG5, EGFR, FANCG, FOS, GFAP, GOLGA2/GM130, GRASP1, UBE2A/hHR6A, HIST1H1 proteins/histone H1, HMGA1, HIVEP3/KRC, UHRF1, KAT5, LMNA, LMNB, LBR, MKI67, LATS1, MAP1B, MAP4, MARCKS, MCM2, MCM4, MKLP1, MLST8, MYB, NEFH, NFIC, NPC/nuclear pore complex, PITPNM1/NIR2, NPM1, NCL, NUCKS1, NPM1/numatrin, ORC1, PRKAR2A, EEF1E1/p18, EIF3F/p47, p53/TP53, NONO/p54NRB, PAPOLA, PLEC/plectin, RB1, TPPP, UL40/R2, RAB4A, RAP1GAP, RBBP8/CtIP, RCC1, RPS6KB1/S6K1, KHDRBS1/SAM68, ESPL1, SKI, BIRC5/survivin, STIP1, TEX14, beta-tubulins, MAPT/TAU, NEDD1, VIM/vimentin, TK1, FOXO1, RUNX1/AML1, SAMHD1, SIRT2, CGAS, GLI2, ZAR1 and RUNX2 (PubMed:10542199, PubMed:19821535). CDK1/CDC2-cyclin-B controls pronuclear union in interphase fertilized eggs (By similarity). Essential for early stages of embryonic development (By similarity). During G2 and early mitosis, CDC25A/B/C-mediated dephosphorylation activates CDK1/cyclin complexes which phosphorylate several substrates that trigger at least centrosome separation, Golgi dynamics, nuclear envelope breakdown and chromosome condensation (By similarity). Once chromosomes are condensed and aligned at the metaphase plate, CDK1 activity is switched off by WEE1- and PKMYT1-mediated phosphorylation to allow sister chromatid separation, chromosome decondensation, reformation of the nuclear envelope and cytokinesis (By similarity). Phosphorylates KRT5 during prometaphase and metaphase (By similarity). Activated by the CDK-activating kinase (CAK) complex consisting of CDK7, cyclin-H/CCNH and MAT1, which is a master regulator of CDK activity (By similarity). Inactivated by PKR/EIF2AK2- and WEE1-mediated phosphorylation upon DNA damage to stop cell cycle and genome replication at the G2 checkpoint thus facilitating DNA repair (By similarity). Reactivated after successful DNA repair through WIP1-dependent signaling leading to CDC25A/B/C- mediated dephosphorylation and restoring cell cycle progression (By similarity). Catalyzes lamin (LMNA, LMNB1 and LMNB2) phosphorylation at the onset of mitosis, promoting nuclear envelope |

| | |
|---|---|
| | breakdown (By similarity). In proliferating cells, CDK1-mediated FOXO1 phosphorylation at the G2-M phase represses FOXO1 interaction with 14- 3-3 proteins and thereby promotes FOXO1 nuclear accumulation and transcription factor activity, leading to cell death of postmitotic neurons (By similarity). The phosphorylation of beta-tubulins regulates microtubule dynamics during mitosis (By similarity). NEDD1 phosphorylation promotes PLK1-mediated NEDD1 phosphorylation and subsequent targeting of the gamma-tubulin ring complex (gTuRC) to the centrosome, an important step for spindle formation (By similarity). In addition, CC2D1A phosphorylation regulates CC2D1A spindle pole localization and association with SCC1/RAD21 and centriole cohesion during mitosis (By similarity). The phosphorylation of Bcl-xL/BCL2L1 after prolongated G2 arrest upon DNA damage triggers apoptosis (By similarity). In contrast, CASP8 phosphorylation during mitosis prevents its activation by proteolysis and subsequent apoptosis (By similarity). This phosphorylation occurs in cancer cell lines, as well as in primary breast tissues and lymphocytes (By similarity). EZH2 phosphorylation promotes H3K27me3 maintenance and epigenetic gene silencing (By similarity). CALD1 phosphorylation promotes Schwann cell migration during peripheral nerve regeneration (PubMed:17200138). CDK1-cyclin-B complex phosphorylates NCKAP5L and mediates its dissociation from centrosomes during mitosis (By similarity). Regulates the amplitude of the cyclic expression of the core clock gene BMAL1 by phosphorylating its transcriptional repressor NR1D1, and this phosphorylation is necessary for SCF(FBXW7)-mediated ubiquitination and proteasomal degradation of NR1D1 (By similarity). Phosphorylates EML3 at 'Thr-881' which is essential for its interaction with HAUS augmin-like complex and TUBG1 (By similarity). Phosphorylates CGAS during mitosis, leading to its inhibition, thereby preventing CGAS activation by self DNA during mitosis (By similarity). Phosphorylates SKA3 during mitosis which promotes SKA3 binding to the NDC80 complex and anchoring of the SKA complex to kinetochores, to enable stable attachment of mitotic spindle microtubules to kinetochores (By similarity). |
| **Cellular Location** | Nucleus. Cytoplasm {ECO:0000250\|UniProtKB:P11440} Mitochondrion {ECO:0000250\|UniProtKB:P11440}. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome {ECO:0000250\|UniProtKB:P06493}. Cytoplasm, cytoskeleton, spindle {ECO:0000250\|UniProtKB:P06493}. Note=Colocalizes with SIRT2 on centrosome during prophase and on splindle fibers during metaphase of the mitotic cell cycle (By similarity). Cytoplasmic during the interphase. Reversibly translocated from cytoplasm to nucleus when phosphorylated before G2-M transition when associated with cyclin-B1 Accumulates in mitochondria in G2-arrested cells upon DNA-damage {ECO:0000250\|UniProtKB:P06493} |

## Background

P34(cdc2), also known as cell division cycle(CDC2), or cyclin-dependent kinase 1(CDK1). CDC2 is a catalytic subunit of a protein kinase complex, called the M-phase promoting factor, that induces entry into mitosis and is universal among eukaryotes. In HeLa cells CDC2 is the most abundant phosphotyrosine-containing protein and its phosphotyrosine content is subject to cell cycle regulation. CDC2 gene is located on chromosome 10.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.