

Anti-GJB2 Antibody

Catalog # ABO10518

Product Information

Application	WB
Primary Accession	P29033
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for Gap junction beta-2 protein(GJB2) detection. Tested with WB in Human;Mouse;Rat.

Reconstitution Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	2706
Other Names	Gap junction beta-2 protein, Connexin-26, Cx26, GJB2
Calculated MW	26215
Application Details	Western blot, 0.1-0.5 µg/ml, Human, Rat, Mouse
Subcellular Localization	Cell membrane; Multi-pass membrane protein. Cell junction, gap junction.
Source	Eukaryota
Protein Name	Gap junction beta-2 protein
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg Thimerosal, 0.05mg NaN3.
Immunogen	A synthetic peptide corresponding to a sequence at the N-terminus of human GJB2(1-16aa, MDWGTLQTILGGVNKH), different from the related mouse and rat sequences by two amino acids.
Purification	Immunogen affinity purified.
Cross Reactivity	No cross reactivity with other proteins
Storage	At -20 °C for one year. After reconstitution, at 4 °C for one month. It can also be aliquotted and stored frozen at -20 °C for a longer time. Avoid repeated freezing and thawing.
Sequence Similarities	Belongs to the connexin family. Beta-type (group I) subfamily.


Protein Information

Name	GJB2
Function	Structural component of gap junctions (PubMed: 16849369 , PubMed: 17551008 , PubMed: 19340074 , PubMed: 19384972 , PubMed: 21094651 , PubMed: 26753910). Gap junctions are dodecameric channels that connect the cytoplasm of adjoining cells. They are formed by the docking of two hexameric hemichannels, one from each cell membrane (PubMed: 17551008 , PubMed: 19340074 , PubMed: 21094651 , PubMed: 26753910). Small molecules and ions diffuse from one cell to a neighboring cell via the central pore (PubMed: 16849369 , PubMed: 19384972 , PubMed: 21094651).
Cellular Location	Cell membrane; Multi-pass membrane protein. Cell junction, gap junction. Note=Colocalizes with GJB4 at gap junction plaques in the cochlea. {ECO:0000250 UniProtKB:Q00977}

Background

Connexin26(CX26), also known as GAP junction protein, beta2, GJB2. Gap junctions were first characterized by electron microscopy as regionally specialized structures on plasma membranes of contacting adherent cells. These structures were shown to consist of cell-to-cell channels. Proteins, called connexins, purified from fractions of enriched gap junctions from different tissues differ. The 3-prime untranslated region of the CX26 transcript contains a putative mRNA instability sequence. The deduced 226-amino acid protein has a calculated molecular mass of about 26 kD. CX26 shares 92.5% identity with rat Cx26. connexin 26(GJB2) is assigned to human chromosome 13q11-q12 .Connexin 26 regulates epidermal barrier and wound remodeling and promotes psoriasiform response. Connexin 26 gene(GJB2) mutation modulates the severity of hearing loss associated with the 1555A-G mitochondrial mutation.

Images



Anti-GJB2 antibody, ABO10518, Western blottingWB: Rat Liver Tissue Lysate

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.