

Anti-GST3/GST Pi Antibody

Catalog # ABO10527

Product Information

Application	WB
Primary Accession	P09211
Host	Rabbit
Reactivity	Human, Mouse
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for Glutathione S-transferase P(GSTP1) detection. Tested with WB in Human;Mouse.

Reconstitution Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	2950
Other Names	Glutathione S-transferase P, 2.5.1.18, GST class-pi, GSTP1-1, GSTP1, FAEE53, GST3
Calculated MW	23356
Application Details	Western blot, 0.1-0.5 µg/ml, Human, Mouse
Subcellular Localization	Cytoplasm . Mitochondrion . Nucleus . The 83 N-terminal amino acids function as un uncleaved transit peptide, and arginine residues within it are crucial for mitochondrial localization.
Source	Eukaryota
Protein Name	Glutathione S-transferase P
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg Thimerosal, 0.05mg NaN ₃ .
Immunogen	A synthetic peptide corresponding to a sequence at the C-terminus of human GST3(197-210aa PEYVNLPINGNGKQ), different from the related mouse sequence by two amino acids.
Purification	Immunogen affinity purified.
Cross Reactivity	No cross reactivity with other proteins
Storage	At -20 °C for one year. After reconstitution, at 4 °C for one month. It can also be aliquotted and stored frozen at -20 °C for a longer time.Avoid repeated

Sequence Similarities

freezing and thawing.
Belongs to the GST superfamily. Pi family.


Protein Information

Name	GSTP1 (HGNC:4638)
Synonyms	FAEES3, GST3
Function	Catalyzes conjugation of reduced glutathione to a wide number of exogenous and endogenous hydrophobic electrophiles (PubMed: 1540159 , PubMed: 1567427 , PubMed: 8433974). Involved in the formation of glutathione conjugates of both prostaglandin A2 (PGA2) and prostaglandin J2 (PGJ2) (PubMed: 9084911). Participates in the formation of novel hepoxilin regioisomers (PubMed: 21046276). Acts as a negative regulator of ferroptosis by mediating glutathione conjugation and detoxification of 4-hydroxynonenal (4-HNE) reactive aldehyde (PubMed: 38016474). Negatively regulates CDK5 activity via p25/p35 translocation to prevent neurodegeneration (PubMed: 21668448).
Cellular Location	Cytoplasm. Mitochondrion. Nucleus. Note=The 83 N-terminal amino acids function as an uncleaved transit peptide, and arginine residues within it are crucial for mitochondrial localization

Background

Glutathione S-transferases pi, also known as GST3, present in all tissues and cells, with the exception of red cells, in which only erythrocyte GST(GSTe) is observed. The GST-pi gene has 7 exons and 6 introns contained within approximately 2.8 kilobases. The GST-pi gene is mapped to chromosome 11. Placental glutathione-S-transferase-pi mRNA is abundantly expressed in human skin. GSTP does not contribute in vivo to the formation of glutathione conjugates of acetaminophen but plays a novel and unexpected role in the toxicity of this compound.

Images



Anti-GST3/GST pi antibody, ABO10527, Western blotting
Lane 1: MCF-7 Cell Lysate
Lane 2: COLO320 Cell Lysate

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.