

Anti-KCNN4 Antibody

Catalog # ABO10533

Product Information

Application	WB
Primary Accession	O15554
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for Intermediate conductance calcium-activated potassium channel protein 4(KCNN4) detection. Tested with WB in Human;Mouse;Rat.
Reconstitution	Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	3783
Other Names	Intermediate conductance calcium-activated potassium channel protein 4, SK4, SKCa 4, SKCa4, IKCa1, IK1, KCa3.1, KCa4, Putative Gardos channel, KCNN4, IK1, IKCA1, KCA4, SK4
Calculated MW	47696
Application Details	Western blot, 0.1-0.5 µg/ml, Human, Rat, Mouse
Subcellular Localization	Membrane; Multi-pass membrane protein.
Tissue Specificity	Widely expressed in non-excitabile tissues.
Source	Eukaryota
Protein Name	Intermediate conductance calcium-activated potassium channel protein 4
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg Thimerosal, 0.05mg Na ₃ N.
Immunogen	A synthetic peptide corresponding to a sequence at the N-terminus of human KCNN4(14-29aa RRKRLLEQEKSLAGWA), different from the related mouse and rat sequences by two amino acids.
Purification	Immunogen affinity purified.
Cross Reactivity	No cross reactivity with other proteins

Storage	At -20° C for one year. After r° Constitution, at 4° C for one month. It° Can also be aliquotted and stored frozen at -20° C for a longer time.Avoid repeated freezing and thawing.
----------------	---

Protein Information

Name	KCNN4 (HGNC:6293)
Synonyms	IK1, IKCA1, KCA4, SK4
Function	Intermediate conductance calcium-activated potassium channel that mediates the voltage-independent transmembrane transfer of potassium across the cell membrane through a constitutive interaction with calmodulin which binds the intracellular calcium allowing its opening (PubMed:10026195, PubMed:10961988, PubMed:11425865, PubMed:15831468, PubMed:17157250, PubMed:18796614, PubMed:26148990, PubMed:9326665, PubMed:9380751, PubMed:9407042). The current is characterized by a voltage-independent activation, an intracellular calcium concentration increase-dependent activation and a single- channel conductance of about 25 picosiemens (PubMed:9326665, PubMed:9380751, PubMed:9407042). Also presents an inwardly rectifying current, thus reducing its already small outward conductance of potassium ions, which is particularly the case when the membrane potential displays positive values, above + 20 mV (PubMed:9326665, PubMed:9380751, PubMed:9407042). Controls calcium influx during vascular contractility by being responsible of membrane hyperpolarization induced by vasoactive factors in proliferative vascular smooth muscle cell types (By similarity). Following calcium influx, the consecutive activation of KCNN4 channel leads to a hyperpolarization of the cell membrane potential and hence an increase of the electrical driving force for further calcium influx promoting sustained calcium entry in response to stimulation with chemotactic peptides (PubMed:26418693). Required for maximal calcium influx and proliferation during the reactivation of naive T-cells (PubMed:17157250, PubMed:18796614). Plays a role in the late stages of EGF-induced macropinocytosis through activation by PI(3)P (PubMed:24591580).
Cellular Location	Cell membrane; Multi-pass membrane protein. Cell projection, ruffle membrane. Note=Targeted to membrane ruffles after EGF stimulation.
Tissue Location	Widely expressed in non-excitabile tissues.

Background

Intermediate conductance calcium-activated potassium channel protein 1(KCNN4, Kca3.1) is part of a potentially heterotetrameric voltage-independent potassium channel that is activated by intracellular calcium. Activation is followed by membrane hyperpolarization, which promotes calcium influx. KCNN4 may be part of the predominant calcium-activated potassium channel in T-lymphocytes. This gene is similar to other KCNN family potassium channel genes, but it differs enough to possibly be considered as part of a new subfamily.

Images

Anti-KCNN antibody, ABO10533, Western blottingAll lanes: Anti KCNN(ABO10533) at 0.5ug/mlWB: HUT Whole Cell Lysate at 40ugPredicted bind size: 60KDObserved bind size: 60KD



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.