

Anti-Topoisomerase II Alpha Antibody

Catalog # ABO10582

Product Information

Application	WB
Primary Accession	P11388
Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for DNA topoisomerase 2-alpha(TOP2A) detection. Tested with WB in Human.

Reconstitution Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	7153
Other Names	DNA topoisomerase 2-alpha, 5.99.1.3, DNA topoisomerase II, alpha isozyme, TOP2A, TOP2
Calculated MW	174385
Application Details	Western blot, 0.1-0.5 µg/ml, Human
Subcellular Localization	Cytoplasm . Nucleus, nucleoplasm . Generally located in the nucleoplasm.
Source	Eukaryota
Protein Name	DNA topoisomerase 2-alpha
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg Thimerosal, 0.05mg NaN ₃ .
Immunogen	A synthetic peptide corresponding to a sequence at the C-terminus of human Topoisomerase II alpha(1466-1480aa RKPSTSDDSDSNFEK), different from the related rat sequence by three amino acids, and different from the related mouse sequence by four amino acids
Purification	Immunogen affinity purified.
Cross Reactivity	No cross reactivity with other proteins
Storage	At -20 °C for one year. After reconstitution, at 4 °C for one month. It can also be aliquotted and stored frozen at -20 °C for a longer time. Avoid repeated freezing and thawing.

Protein Information

Name	TOP2A
Synonyms	TOP2
Function	Key decatenating enzyme that alters DNA topology by binding to two double-stranded DNA molecules, generating a double-stranded break in one of the strands, passing the intact strand through the broken strand, and religating the broken strand (PubMed: 17567603 , PubMed: 18790802 , PubMed: 22013166 , PubMed: 22323612 , PubMed: 27754753). May play a role in regulating the period length of BMAL1 transcriptional oscillation (By similarity).
Cellular Location	Cytoplasm. Nucleus, nucleoplasm. Nucleus. Nucleus, nucleolus
Tissue Location	Expressed in the tonsil, spleen, lymph node, thymus, skin, pancreas, testis, colon, kidney, liver, brain and lung (PubMed:9155056). Also found in high-grade lymphomas, squamous cell lung tumors and seminomas (PubMed:9155056)

Background

The human topoisomerase II enzyme is encoded by a single-copy gene which is mapped to 17q21-q22. The TOP2A gene spans approximately 30 kb and contains 35 exons. Furthermore, DNA topoisomerases are enzymes that control and alter the topologic states of DNA in both prokaryotes and eukaryotes. Topoisomerase II from eukaryotic cells catalyzes the relaxation of supercoiled DNA molecules, catenation, decatenation, knotting, and unknotting of circular DNA. It appears likely that the reaction catalyzed by topoisomerase II involves the crossing-over of 2 DNA segments. There are about 100,000 molecules of topoisomerase II per HeLa cell nucleus, constituting about 0.1% of the nuclear extract. DNA topoisomerase II-alpha is associated with the pol II holoenzyme and is a required component of chromatin-dependent coactivation. Specific inhibitors of topoisomerase II blocked transcription on chromatin templates, but did not affect transcription on naked templates. Addition of purified topoisomerase II-alpha reconstituted chromatin-dependent activation activity in reactions with core pol II.

Images



Anti- Topoisomerase II alpha antibody, ABO10582,
Western blottingAll lanes: Anti Topoisomerase II alpha
(ABO10582) at 0.5ug/mlLane 1: HELA Whole Cell Lysate at
40ugLane 2: JURKAT Whole Cell Lysate at 40ugPredicted
bind size: 174KDObserved bind size: 174KD