

Anti-Chk2 Antibody

Catalog # ABO10587

Product Information

Application	WB
Primary Accession	O96017
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for Serine/threonine-protein kinase Chk2(CHEK2) detection. Tested with WB in Human;Mouse;Rat.
Reconstitution	Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	11200
Other Names	Serine/threonine-protein kinase Chk2, 2.7.11.1, CHK2 checkpoint homolog, Cds1 homolog, Hucds1, hCds1, Checkpoint kinase 2, CHEK2, CDS1, CHK2, RAD53
Calculated MW	60915
Application Details	Western blot, 0.1-0.5 µg/ml, Human, Rat, Mouse
Subcellular Localization	Isoform 2: Nucleus. Isoform 10 is present throughout the cell.
Tissue Specificity	High expression is found in testis, spleen, colon and peripheral blood leukocytes. Low expression is found in other tissues.
Source	Eukaryota
Protein Name	Serine/threonine-protein kinase Chk2
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg Thimerosal, 0.05mg Na ₃ N.
Immunogen	A synthetic peptide corresponding to a sequence at the C-terminus of human Chk2(427-441aa FSEHRTQVSLKDQIT), different from the related rat and mouse sequences by one amino acid.
Purification	Immunogen affinity purified.
Cross Reactivity	No cross reactivity with other proteins

Storage	At -20° C for one year. After r° Constitution, at 4° C for one month. It° Can also be aliquotted and stored frozen at -20° C for a longer time.Avoid repeated freezing and thawing.
Sequence Similarities	Belongs to the protein kinase superfamily. CAMK Ser/Thr protein kinase family. CHK2 subfamily.

Protein Information


Name	CHEK2 (HGNC:16627)
Synonyms	CDS1, CHK2, RAD53
Function	Serine/threonine-protein kinase which is required for checkpoint-mediated cell cycle arrest, activation of DNA repair and apoptosis in response to the presence of DNA double-strand breaks. May also negatively regulate cell cycle progression during unperturbed cell cycles. Following activation, phosphorylates numerous effectors preferentially at the consensus sequence [L-X-R-X-X-S/T] (PubMed: 37943659). Regulates cell cycle checkpoint arrest through phosphorylation of CDC25A, CDC25B and CDC25C, inhibiting their activity. Inhibition of CDC25 phosphatase activity leads to increased inhibitory tyrosine phosphorylation of CDK-cyclin complexes and blocks cell cycle progression. May also phosphorylate NEK6 which is involved in G2/M cell cycle arrest. Regulates DNA repair through phosphorylation of BRCA2, enhancing the association of RAD51 with chromatin which promotes DNA repair by homologous recombination. Also stimulates the transcription of genes involved in DNA repair (including BRCA2) through the phosphorylation and activation of the transcription factor FOXM1. Regulates apoptosis through the phosphorylation of p53/TP53, MDM4 and PML. Phosphorylation of p53/TP53 at 'Ser-20' by CHEK2 may alleviate inhibition by MDM2, leading to accumulation of active p53/TP53. Phosphorylation of MDM4 may also reduce degradation of p53/TP53. Also controls the transcription of pro-apoptotic genes through phosphorylation of the transcription factor E2F1. Tumor suppressor, it may also have a DNA damage-independent function in mitotic spindle assembly by phosphorylating BRCA1. Its absence may be a cause of the chromosomal instability observed in some cancer cells. Promotes the CCAR2-SIRT1 association and is required for CCAR2-mediated SIRT1 inhibition (PubMed: 25361978). Under oxidative stress, promotes ATG7 ubiquitination by phosphorylating the E3 ubiquitin ligase TRIM32 at 'Ser-55' leading to positive regulation of the autophagosome assembly (PubMed: 37943659).
Cellular Location	[Isoform 2]: Nucleus. Note=Isoform 10 is present throughout the cell [Isoform 7]: Nucleus. [Isoform 12]: Nucleus.
Tissue Location	High expression is found in testis, spleen, colon and peripheral blood leukocytes. Low expression is found in other tissues

Background

CHEK2, a protein kinase that is activated in response to DNA damage, is involved in cell cycle arrest. Mapped on 22q12.1, CHK2 has a potential regulatory region rich in SQ and TQ amino acid pairs. It regulates BRCA1 function after DNA damage by phosphorylating serine-988 of BRCA1. Additionally, CHK2 can be modified by phosphorylation and activated in response to ionizing radiation, and can be also modified in response to hydroxyurea treatment. Furthermore, oligomerization of CHEK2 increases the efficiency of transautophosphorylation, resulting in the release of active CHEK2 monomers that proceed to enforce checkpoint control in irradiated cells. Moreover, CHK2 is a tumor suppressor gene conferring predisposition to sarcoma, breast cancer, and brain tumors, and that their observations provided a link between the central role of p53 inactivation in human cancer and the well-defined G2 checkpoint in yeast. There is a wide

expression of small amounts of CHK2 mRNA with larger amounts in human testis, spleen, colon, and peripheral blood leukocytes.

Images



Anti-Chk2 antibody, ABO10587, Western blottingWB:
MCF-7 Cell Lysate

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.