

Anti-uPAR Antibody

Catalog # ABO10669

Product Information

Application	WB, IHC-P
Primary Accession	Q03405
Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for Urokinase plasminogen activator surface receptor(PLAUR) detection. Tested with WB, IHC-P in Human.
Reconstitution	Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	5329
Other Names	Urokinase plasminogen activator surface receptor, U-PAR, uPAR, Monocyte activation antigen Mo3, CD87, PLAUR, MO3, UPAR
Calculated MW	36978
Application Details	Immunohistochemistry(Paraffin-embedded Section), 0.5-1 µg/ml, Human, By Heat Western blot, 0.1-0.5 µg/ml, Human
Subcellular Localization	Cell membrane . Cell projection, invadopodium membrane . Colocalized with FAP (seprase) preferentially at the cell surface of invadopodia membrane in a cytoskeleton-, integrin- and vitronectin-dependent manner. .
Tissue Specificity	Expressed in neurons of the rolandic area of the brain (at protein level). Expressed in the brain.
Source	Eukaryota
Protein Name	Urokinase plasminogen activator surface receptor(U-PAR/uPAR)
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg Thimerosal, 0.05mg Na ₃ N.
Immunogen	A synthetic peptide corresponding to a sequence at the C-terminus of human uPA Receptor(293-304aa CNHPDLVDVQYRS).
Purification	Immunogen affinity purified.

Cross Reactivity	No cross reactivity with other proteins
Storage	At -20 °C for one year. After r ° Constitution, at 4 °C for one month. It ° Can also be aliquotted and stored frozen at -20 °C for a longer time.Avoid repeated freezing and thawing.
Sequence Similarities	Contains 3 UPAR/Ly6 domains.

Protein Information



Name	PLAUR (HGNC:9053)
Synonyms	MO3, UPAR
Function	GPI-anchored receptor that recruits and activates the urokinase-type plasminogen activator (uPA) at the cell surface. Activated uPA converts plasminogen into plasmin, initiating extracellular matrix degradation and remodeling (PubMed: 1689240 , PubMed: 15677461). Also binds the extracellular matrix protein vitronectin, promoting cell-matrix adhesion and indirectly integrin signaling (PubMed: 17548516 , PubMed: 25168639 , PubMed: 28849762). This dual interaction coordinates dynamic changes in cell migration, proliferation, and tissue remodeling (PubMed: 28849762).
Cellular Location	[Isoform 1]: Cell membrane; Lipid-anchor, GPI-anchor. Cell projection, invadopodium membrane; Lipid-anchor, GPI-anchor. Note=Colocalized with FAP (seprase) preferentially at the cell surface of invadopodia membrane in a cytoskeleton-, integrin- and vitronectin-dependent manner
Tissue Location	Expressed in neurons of the rolandic area of the brain (at protein level). Expressed in the brain

Background

PLAUR(PLASMINOGEN ACTIVATOR RECEPTOR, UROKINASE-TYPE), also known as UPAR or CD87, is multidomain glycoprotein tethered to the cell membrane with a glycosylphosphatidylinositol(GPI) anchor. PLAUR consists of three different domains of the Ly-6/uPAR/alpha-neurotoxin family. PLAUR is originally identified as a saturable binding site for urokinase on the cell surface. And the gene plays an important role in many normal as well as pathologic processes. The PLAUR gene is localized to 19q13.31. PLAUR is a part of the plasminogen activation system, which in the healthy body is involved in tissue reorganization events such as mammary gland involution and wound healing. PLAUR binds urokinase and thus restricts plasminogen activation to the immediate vicinity of the cell membrane. Thus it seems to be an important player in the regulation of this process. In human coronary artery vascular smooth muscle cells, UPA stimulates cell migration via a UPAR signaling complex containing TYK2 and phosphatidylinositol 3-kinase.

Images

Anti-uPA Receptor antibody, ABO10669, Western blotting
Lane 1: MCF-7 Cell Lysate Lane 2: HELA Cell Lysate
Lane 3: RAJI Cell Lysate Lane 4: SMMC Cell Lysate



Anti-uPA Receptor antibody, ABO10669, IHC(P)IHC(P):
Human Mammary Cancer Tissue

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.