10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Anti-MMP-9 Antibody

Catalog # ABO10680

Product Information

Application	WB
Primary Accession	P14780
Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for Matrix metalloproteinase-9(MMP9) detection. Tested with WB in Human.
Reconstitution	Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	4318
Other Names	Matrix metalloproteinase-9, MMP-9, 3.4.24.35, 92 kDa gelatinase, 92 kDa type IV collagenase, Gelatinase B, GELB, 67 kDa matrix metalloproteinase-9, 82 kDa matrix metalloproteinase-9, MMP9, CLG4B
Calculated MW	78458
Application Details	Western blot, 0.1-0.5 □g/ml, Human, Rat
Subcellular Localization	Secreted, extracellular space, extracellular matrix .
Tissue Specificity	Produced by normal alveolar macrophages and granulocytes.
Source	Eukaryota
Protein Name	Matrix metalloproteinase-9(MMP-9)
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg Thimerosal, 0.05mg NaN3.
Immunogen	A synthetic peptide corresponding to a sequence at the C-terminus of human MMP-9(689-705aa NQVDQVGYVTYDILQCP), different from the related mouse sequence by two amino acids, and from the related rat sequence by three amino acids.
Purification	Immunogen affinity purified.
Cross Reactivity	No cross reactivity with other proteins

Storage	At -20˚C for one year. After r˚Constitution, at 4˚C for one month. It˚Can also be aliquotted and stored frozen at -20˚C for a longer time.Avoid repeated freezing and thawing.
Sequence Similarities	Belongs to the peptidase M10A family.


Protein Information

Name	MMP9
Synonyms	CLG4B
Function	Matrix metalloproteinase that plays an essential role in local proteolysis of the extracellular matrix and in leukocyte migration (PubMed:12879005, PubMed:1480034, PubMed:2551898). Could play a role in bone osteoclastic resorption (By similarity). Cleaves KiSS1 at a Gly-|-Leu bond (PubMed:12879005). Cleaves NINJ1 to generate the Secreted ninjurin-1 form (PubMed:32883094). Cleaves type IV and type V collagen into large C-terminal three quarter fragments and shorter N- terminal one quarter fragments (PubMed:1480034). Degrades fibronectin but not laminin or Pz-peptide.
Cellular Location	Secreted, extracellular space, extracellular matrix
Tissue Location	Detected in neutrophils (at protein level) (PubMed:7683678). Produced by normal alveolar macrophages and granulocytes.

Background

Matrix metallopeptidase 9(MMP-9), also known as 92 kDa type IV collagenase, 92 kDa gelatinase or gelatinase B(GELB), is an enzyme that in humans is encoded by the MMP9 gene. Proteins of the matrix metalloproteinase(MMP) family are involved in the breakdown of extracellular matrix in normal physiological processes. Most MMPs are secreted as inactive proproteins which are activated when cleaved by extracellular proteinases. The enzyme encoded by this gene degrades type IV and V collagens. Studies in rhesus monkeys suggest that the enzyme is involved in IL-8-induced mobilization of hematopoietic progenitor cells from bone marrow, and murine studies suggest a role in tumor-associated tissue remodeling.

Images



Anti-MMP-9 antibody, ABO10680, Western blottingLane 1: Rat Embryo Tissue LysateLane 2: MM453 Cell LysateLane 3: HELA Cell LysateLane 4: SMMC Cell LysateLane 5: JURKAT Cell LysateLane 6: HT1080 Cell Lysate

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.