

Anti-CXCL10/IP-10 Antibody

Catalog # ABO10719

Product Information

Application	WB, IHC-P
Primary Accession	P02778
Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for C-X-C motif chemokine 10(CXCL10) detection. Tested with WB, IHC-P in Human.

Reconstitution Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	3627
Other Names	C-X-C motif chemokine 10, 10 kDa interferon gamma-induced protein, Gamma-IP10, IP-10, Small-inducible cytokine B10, CXCL10(1-73), CXCL10, INP10, SCYB10
Calculated MW	10881
Application Details	Immunohistochemistry(Paraffin-embedded Section), 0.5-1 µg/ml, Human, By Heat Western blot, 0.1-0.5 µg/ml, Human
Subcellular Localization	Secreted.
Source	Eukaryota
Protein Name	C-X-C motif chemokine 10
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg Thimerosal, 0.05mg NaN ₃ .
Immunogen	A synthetic peptide corresponding to a sequence at the C-terminus of human IP10(79-98aa SKAIKNLLKAVSKERSKRSP).
Purification	Immunogen affinity purified.
Cross Reactivity	No cross reactivity with other proteins
Storage	At -20° C for one year. After reconstitution, at 4° C for one month. It can also

	be aliquotted and stored frozen at -20° C for a longer time.Avoid repeated freezing and thawing.
Sequence Similarities	Belongs to the intercrine alpha (chemokine CxC) family.

Protein Information



Name	CXCL10
Synonyms	INP10, SCYB10
Function	Pro-inflammatory cytokine that is involved in a wide variety of processes such as chemotaxis, differentiation, and activation of peripheral immune cells, regulation of cell growth, apoptosis and modulation of angiostatic effects (PubMed: 11157474 , PubMed: 22652417 , PubMed: 7540647). Plays thereby an important role during viral infections by stimulating the activation and migration of immune cells to the infected sites (By similarity). Mechanistically, binding of CXCL10 to the CXCR3 receptor activates G protein-mediated signaling and results in downstream activation of phospholipase C-dependent pathway, an increase in intracellular calcium production and actin reorganization (PubMed: 12750173 , PubMed: 19151743). In turn, recruitment of activated Th1 lymphocytes occurs at sites of inflammation (PubMed: 12663757 , PubMed: 12750173). Activation of the CXCL10/CXCR3 axis also plays an important role in neurons in response to brain injury for activating microglia, the resident macrophage population of the central nervous system, and directing them to the lesion site. This recruitment is an essential element for neuronal reorganization (By similarity).
Cellular Location	Secreted.
Tissue Location	Mainly secreted by monocytes, endothelial cells as well as fibroblasts. Expressed by epithelial cells in thymus (PubMed:11157474). Microglial cells produce CXCL10 in response to viral stimulation (PubMed:12663757).

Background

C-X-C motif chemokine 10(CXCL10), also known as Interferon gamma-induced protein 10 kDa(IP-10) or small-inducible cytokine B10, is a protein that in humans is encoded by the CXCL10 gene. It is a small cytokine belonging to the CXC chemokine family. The gene for CXCL10 is located on human chromosome 4 in a cluster among several other CXC chemokines. Luster et al.(1987) reported the isolation of an interferon-inducible gene that encodes a 98-amino acid protein called IP10. The chemokine IFN-gamma-inducible protein 10(IP-10; CXCL10), a CXC chemokine, and its receptor, CXCR3, were found to be overexpressed in lymph node cells of EAMG rats. IP10 inhibits bone marrow colony formation, has antitumor activity in vivo, is a chemoattractant for human monocytes and T cells, and promotes T cell adhesion to endothelial cells.

Images

Anti-IP10 antibody, ABO10719, Western blotting
Lane 1: Recombinant Human CXCL10 Protein 10ng
Lane 2: Recombinant Human CXCL10 Protein 5ng
Lane 3: Recombinant Human CXCL10 Protein 2.5ng



Anti-IP10 antibody, ABO10719, IHC(P)IHC(P): Human Mammary Cancer Tissue

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.