

Anti-IL-7 Antibody

Catalog # ABO10786

Product Information

Application	WB, IHC-P
Primary Accession	P10168
Host	Rabbit
Reactivity	Mouse
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for Interleukin-7(IL7) detection. Tested with WB, IHC-P in Mouse.

Reconstitution Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	16196
Other Names	Interleukin-7, IL-7, Il7, IL-7
Calculated MW	17727
Application Details	Immunohistochemistry(Paraffin-embedded Section), 0.5-1 µg/ml, Mouse, By Heat Western blot, 0.1-0.5 µg/ml, Mouse
Subcellular Localization	Secreted.
Source	Eukaryota
Protein Name	Interleukin-7(IL-7)
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg Thimerosal, 0.05mg NaN3.
Immunogen	A synthetic peptide corresponding to a sequence at the N-terminus of mouse IL7(26-42aa ECHIKDKEGKAYESVLM).
Purification	Immunogen affinity purified.
Cross Reactivity	No cross reactivity with other proteins
Storage	At -20° C for one year. After r° Constitution, at 4° C for one month. It° Can also be aliquotted and stored frozen at -20° C for a longer time.Avoid repeated freezing and thawing.


Protein Information

Name	Il7
Synonyms	Il-7
Function	Hematopoietic cytokine that plays an essential role in the development, expansion, and survival of naive and memory T-cells and B- cells thereby regulating the number of mature lymphocytes and maintaining lymphoid homeostasis (PubMed: 28811625 , PubMed: 7699333). Mechanistically, exerts its biological effects through a receptor composed of IL7RA subunit and the cytokine receptor common subunit gamma/CSF2RG. Binding to the receptor leads to activation of various kinases including JAK1 or JAK3 depending on the cell type and subsequently propagation of signals through activation of several downstream signaling pathways including the PI3K/Akt/mTOR or the JAK- STAT5.
Cellular Location	Secreted {ECO:0000250 UniProtKB:P13232}.

Background


IL-7, Interleukin 7, is a protein that in humans is encoded by the IL-7 gene. IL-7 a hematopoietic growth factor secreted by stromal cells in the red marrow and thymus. It is also produced by keratinocytes, dendritic cells, hepatocytes, neurons, and epithelial cells but is not produced by lymphocytes. By the combination of approaches, IL-7 is located on 8q12-q13. IL-7 is critical for early T-cell development and homeostasis of naive and memory CD8-positive T cells. It signals through the IL-7 receptor complex, which consists of the IL7R-alpha chain and the common IL2R-gamma chain is critical for early T-cell development and homeostasis of naive and memory CD8 -positive T cells. It signals through the IL-7 receptor complex, which consists of the IL7R-alpha chain and the common IL2R-gamma chain.

Images



Anti-IL7 antibody, ABO10786, Western blotting
Lane 1:
Recombinant Mouse IL-7 Protein 10ng
Lane 2:
Recombinant Mouse IL-7 Protein 5ng
Lane 3:
Recombinant Mouse IL-7 Protein 2.5ng

Anti-IL7 antibody, ABO10786, IHC(P)
IHC(P): Mouse Spleen Tissue



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.