

Anti-Aquaporin 3 Antibody

Catalog # ABO10807

Product Information

Application	WB, IHC-P, IHC-F, ICC
Primary Accession	Q92482
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for Aquaporin-3(AQP3) detection. Tested with WB, IHC-P, IHC-F, ICC in Human;Mouse;Rat.
Reconstitution	Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	360
Other Names	Aquaporin-3, AQP-3, Aquaglyceroporin-3, AQP3
Calculated MW	31544
Application Details	Immunohistochemistry(Paraffin-embedded Section), 0.5-1 μ g/ml, Human, Rat, Mouse, By Heat Immunocytochemistry , 0.5-1 μ g/ml, Human, Mouse, Rat Immunohistochemistry(Frozen Section), 0.5-1 μ g/ml, Rat, Human, Mouse Western blot, 0.1-0.5 μ g/ml, Human, Mouse, Rat
Subcellular Localization	Basolateral cell membrane; Multi-pass membrane protein. In collecting ducts of kidney.
Tissue Specificity	Widely expressed in epithelial cells of kidney (collecting ducts) and airways, in keratinocytes, immature dendritic cells and erythrocytes. Isoform 2 is not detectable in erythrocytes at the protein level.
Source	Eukaryota
Protein Name	Aquaporin-3
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg Thimerosal, 0.05mg NaN ₃ .
Immunogen	A synthetic peptide corresponding to a sequence at the C-terminus of human Aquaporin 3(278-292aa EENVKLAHVKHKEQI), different from the related rat and mouse sequences by one amino acid.

Purification	Immunogen affinity purified.
Cross Reactivity	No cross reactivity with other proteins
Storage	At -20° C for one year. After r° Constitution, at 4° C for one month. It° Can also be aliquotted and stored frozen at -20° C for a longer time.Avoid repeated freezing and thawing.
Sequence Similarities	Belongs to the MIP/aquaporin (TC 1.A.8) family.

Protein Information



Name	AQP3 {ECO:0000303 PubMed:7558005, ECO:0000312 HGNC:HGNC:636}
Function	Aquaglyceroporins form homotetrameric transmembrane channels, with each monomer independently mediating glycerol and water transport across the plasma membrane along their osmotic gradient (PubMed: 12239222 , PubMed: 30420639). Could also be permeable to urea (By similarity). Also participates in cell permeability to H2O2 and H2O2- mediated signaling (PubMed: 20724658). In skin, transports glycerol to the epidermis and stratum corneum, where it maintains hydration, elasticity, and supports lipid biosynthesis for barrier repair (By similarity). In kidney, contributes to the reabsorption of water, helping the body maintain proper fluid balance (By similarity).
Cellular Location	Cell membrane; Multi-pass membrane protein {ECO:0000250 UniProtKB:O14520}. Basolateral cell membrane {ECO:0000250 UniProtKB:P47862}; Multi-pass membrane protein {ECO:0000250 UniProtKB:O14520}
Tissue Location	Widely expressed in epithelial cells of kidney (collecting ducts) and airways, in keratinocytes, immature dendritic cells and erythrocytes. Isoform 2 is not detectable in erythrocytes at the protein level

Background


This gene encodes the water channel protein aquaporin 3. Aquaporins are a family of small integral membrane proteins related to the major intrinsic protein, also known as aquaporin 0. Aquaporin 3 is localized at the basal lateral membranes of collecting duct cells in the kidney. In addition to its water channel function, aquaporin 3 has been found to facilitate the transport of nonionic small solutes such as urea and glycerol, but to a smaller degree. It has been suggested that water channels can be functionally heterogeneous and possess water and solute permeation mechanisms. Alternative splicing of this gene results in multiple transcript variants encoding different isoforms.

Images


Anti-Aquaporin 3 antibody, ABO10807, Western blottingAll lanes: Anti (ABO10807) at 0.5ug/mlLane 1: Rat Kidney Tissue Lysate at 50ugLane 2: Rat Lung Tissue Lysate at 50ugLane 3: Mouse Kidney Tissue Lysate at 50ugLane 4: MM453 Whole Cell Lysate at 40ugLane 5: SMMC-7721 Whole Cell Lysate at 40ugPredicted bind size: 32KDObserved bind size: 32KD




Anti-Aquaporin 3 antibody, ABO10807, IHC(P)IHC(P): Rat Kidney Tissue



Anti-Aquaporin 3 antibody, ABO10807, ICCICC: HELA Cell




Anti-Aquaporin 3 antibody, ABO10807, IHC(F)IHC(F): Rat Kidney Tissue



Anti-Aquaporin 3 antibody, ABO10807, IHC(P)IHC(P): Human Renal Cancer Tissue

Anti-Aquaporin 3 antibody, ABO10807, IHC(P)IHC(P): Human Lung Cancer Tissue



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.