

Anti-GRB7 Antibody

Catalog # ABO10903

Product Information

Application	WB, IHC-P
Primary Accession	Q14451
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for Growth factor receptor-bound protein 7(GRB7) detection. Tested with WB, IHC-P in Human;Mouse;Rat.
Reconstitution	Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	2886
Other Names	Growth factor receptor-bound protein 7, B47, Epidermal growth factor receptor GRB-7, GRB7 adapter protein, GRB7
Calculated MW	59681
Application Details	Immunohistochemistry(Paraffin-embedded Section), 0.5-1 μ g/ml, Human, Mouse, Rat, By Heat Western blot, 0.1-0.5 μ g/ml, Human, Rat, Mouse
Subcellular Localization	Cytoplasm. Cell junction, focal adhesion. Cell membrane; Peripheral membrane protein; Cytoplasmic side. Cytoplasmic granule . Cell projection. Predominantly cytoplasmic. Detected in stress granules, where mRNA is stored under stress conditions (By similarity). .
Source	Eukaryota
Protein Name	Growth factor receptor-bound protein 7
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg Thimerosal, 0.05mg Na ₃ .
Immunogen	A synthetic peptide corresponding to a sequence in the middle region of human GRB7(124-141aa ARHVCEMLVQRAHALSDE), identical to the related rat and mouse sequences.
Purification	Immunogen affinity purified.
Cross Reactivity	No cross reactivity with other proteins

Storage	At -20 °C for one year. After r ° Constitution, at 4 °C for one month. It ° Can also be aliquotted and stored frozen at -20 °C for a longer time.Avoid repeated freezing and thawing.
Sequence Similarities	Belongs to the GRB7/10/14 family.


Protein Information

Name	GRB7
Function	Adapter protein that interacts with the cytoplasmic domain of numerous receptor kinases and modulates down-stream signaling. Promotes activation of down-stream protein kinases, including STAT3, AKT1, MAPK1 and/or MAPK3. Promotes activation of HRAS. Plays a role in signal transduction in response to EGF. Plays a role in the regulation of cell proliferation and cell migration. Plays a role in the assembly and stability of RNA stress granules. Binds to the 5'UTR of target mRNA molecules and represses translation of target mRNA species, when not phosphorylated. Phosphorylation impairs RNA binding and promotes stress granule disassembly during recovery after cellular stress (By similarity).
Cellular Location	Cytoplasm. Cell junction, focal adhesion. Cell membrane; Peripheral membrane protein; Cytoplasmic side. Cytoplasmic granule {ECO:0000250 UniProtKB:Q03160}. Cell projection. Note=Predominantly cytoplasmic. Detected in stress granules, where mRNA is stored under stress conditions {ECO:0000250 UniProtKB:Q03160}

Background


GRB7, growth factor receptor-bound protein 7, is a protein which in humans is encoded by the GRB7 gene. The product of this gene belongs to a small family of adaptor proteins that are known to interact with a number of receptor tyrosine kinases and signaling molecules. This gene encodes a growth factor receptor-binding protein that interacts with epidermal growth factor receptor (EGFR) and ephrin receptors. The protein plays a role in the integrin signaling pathway and cell migration by binding with focal adhesion kinase (FAK). Alternative splicing results in multiple transcript variants encoding different isoforms, although the full-length natures of only two of the variants have been determined to date. GRB7 is an SH2-domain adaptor protein that binds to receptor tyrosine kinases and provides the intra-cellular direct link to the Ras proto-oncogene. Human GRB7 is located on the long arm of chromosome 17, next to the ERBB2 (alias HER2/neu) proto-oncogene. These two genes are commonly co-amplified (present in excess copies) in breast cancers. GRB7 thought to be involved in migration, is well known to be over-expressed in testicular germ cell tumors, esophageal cancers, and gastric cancers.

Images



Anti-GRB7 antibody, ABO10903, IHC(P)IHC(P): Human Oesophagus Squama Cancer Tissue

Anti-GRB7 antibody, ABO10903, All Western blottingAll lanes: Anti-GRB7(ABO10903) at 0.5ug/mlLane 1: Rat Testis



Tissue Lysate at 40ug
 Lane 2: SMMC Whole Cell Lysate at 40ug
 Lane 3: HELA Whole Cell Lysate at 40ug
 Lane 4: A549 Whole Cell Lysate at 40ug
 Lane 5: SW620 Whole Cell Lysate at 40ug
 Predicted bind size: 60KD
 Observed bind size: 60KD

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.