

Anti-NOX2/gp91phox Antibody

Catalog # ABO10982

Product Information

Application	WB, IHC-P
Primary Accession	P04839
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for Cytochrome b-245 heavy chain(CYBB) detection. Tested with WB, IHC-P in Human;Mouse;Rat.
Reconstitution	Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	1536
Other Names	Cytochrome b-245 heavy chain, 1.-.-., CGD91-phox, Cytochrome b(558) subunit beta, Cytochrome b558 subunit beta, Heme-binding membrane glycoprotein gp91phox, NADPH oxidase 2, Neutrophil cytochrome b 91 kDa polypeptide, Superoxide-generating NADPH oxidase heavy chain subunit, gp91-1, gp91-phox, p22 phagocyte B-cytochrome, CYBB, NOX2
Calculated MW	65336
Application Details	Immunohistochemistry(Paraffin-embedded Section), 0.5-1 µg/ml, Human, Rat, Mouse, By Heat Western blot, 0.1-0.5 µg/ml, Human, Rat, Mouse
Subcellular Localization	Cell membrane; Multi-pass membrane protein.
Tissue Specificity	Detected in neutrophils (at protein level). .
Source	Eukaryota
Protein Name	Cytochrome b-245 heavy chain
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg Thimerosal, 0.05mg NaN3.
Immunogen	A synthetic peptide corresponding to a sequence in the middle region of human NOX2(442-459aa YWLCRDTHAFEWFADLLQ), identical to the related rat and mouse sequences.

Purification	Immunogen affinity purified.
Cross Reactivity	No cross reactivity with other proteins
Storage	At -20° C for one year. After r° Constitution, at 4° C for one month. It° Can also be aliquotted and stored frozen at -20° C for a longer time.Avoid repeated freezing and thawing.
Sequence Similarities	Contains 1 FAD-binding FR-type domain.

Protein Information



Name	CYBB (HGNC:2578)
Synonyms	NOX2
Function	Catalytic subunit of the phagocyte NADPH oxidase complex that mediates the transfer of electrons from cytosolic NADPH to O2 to produce the superoxide anion (O2(-)) (PubMed: 15338276 , PubMed: 36241643 , PubMed: 36413210 , PubMed: 38355798). In the activated complex, electrons are first transferred from NADPH to flavin adenine dinucleotide (FAD) and subsequently transferred via two heme molecules to molecular oxygen, producing superoxide through an outer-sphere reaction (Probable) (PubMed: 38355798). Activation of the NADPH oxidase complex is initiated by the assembly of cytosolic subunits of the NADPH oxidase complex with the core NADPH oxidase complex to form a complex at the plasma membrane or phagosomal membrane (PubMed: 19028840 , PubMed: 38355798). This activation process is initiated by phosphorylation dependent binding of the cytosolic NCF1/p47-phox subunit to the C-terminus of CYBA/p22-phox (By similarity). NADPH oxidase complex assembly is impaired through interaction with NRROS (By similarity).
Cellular Location	Cell membrane; Multi-pass membrane protein. Note=As unassembled monomer may localize to the endoplasmic reticulum
Tissue Location	Detected in neutrophils (at protein level).

Background

NOX2(NADPH OXIDASE 2), also called CYBB(CYTOCHROME b(-245), BETA SUBUNIT), p91-PHOX or GP91-1, is a human gene encoding a glycoprotein.NOX2 is an essential component of phagocytic NADPH-oxidase, a membrane-bound enzyme complex that generates large quantities of microbicidal superoxide and other oxidants upon activation. It is mapped on Xp11.4. NOX2 is a heterodimer composed of an alpha chain of relative molecular mass 23 kD and a beta chain of 76 to 82 kD. NOX2 assembled on DC phagosomes in a gp91-phox subunit-dependent manner, and that reactive oxygen species were produced in a more sustained manner in immature DC phagosomes than in macrophage phagosomes. As a major player in innate immune responses in neutrophils, NOX2 is also involved in adaptive immunity through its activity in DCs. In heart cells, physiologic stretch rapidly activates reduced-form NOX2 to produce reactive oxygen species(ROS) in a process dependent on microtubules(X-ROS signaling).

Images

Anti-NOX2/gp91phox antibody, ABO10982, Western blottingLane 1: HELA Cell Lysate Lane 2: JURKAT Cell LysateLane 3: MCF-7 Cell Lysate Lane 4: SMMC Cell Lysate



Anti-NOX2/gp91phox antibody, ABO10982, IHC(P)IHC(P):
Human Intestinal Cancer Tissue

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.