

Anti-SPTLC1 Antibody

Catalog # ABO11084

Product Information

Application	WB, IHC-P
Primary Accession	O15269
Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for Serine palmitoyltransferase 1(SPTLC1) detection. Tested with WB, IHC-P in Human.
Reconstitution	Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	10558
Other Names	Serine palmitoyltransferase 1, 2.3.1.50, Long chain base biosynthesis protein 1, LCB 1, Serine-palmitoyl-CoA transferase 1, SPT 1, SPT1, SPTLC1, LCB1
Calculated MW	52744
Application Details	Immunohistochemistry(Paraffin-embedded Section), 0.5-1 µg/ml, Human, By Heat Western blot, 0.1-0.5 µg/ml, Human
Subcellular Localization	Endoplasmic reticulum membrane ; Single-pass membrane protein .
Tissue Specificity	Widely expressed. Not detected in small intestine. .
Source	Eukaryota
Protein Name	Serine palmitoyltransferase 1
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg Thimerosal, 0.05mg NaN ₃ .
Immunogen	A synthetic peptide corresponding to a sequence in the middle region of human SPTLC1(138-154aa FYGTFDVHLDLEDRLAK).
Purification	Immunogen affinity purified.
Cross Reactivity	No cross reactivity with other proteins
Storage	At -20 °C for one year. After reconstitution, at 4 °C for one month. It can also

Sequence Similarities	be aliquotted and stored frozen at -20 °C for a longer time.Avoid repeated freezing and thawing. Belongs to the class-II pyridoxal-phosphate-dependent aminotransferase family.
------------------------------	--

Protein Information


Name	SPTLC1
Synonyms	LCB1
Function	Component of the serine palmitoyltransferase multisubunit enzyme (SPT) that catalyzes the initial and rate-limiting step in sphingolipid biosynthesis by condensing L-serine and activated acyl-CoA (most commonly palmitoyl-CoA) to form long-chain bases. The SPT complex is also composed of SPTLC2 or SPTLC3 and SPTSSA or SPTSSB. Within this complex, the heterodimer with SPTLC2 or SPTLC3 forms the catalytic core (PubMed: 19416851 , PubMed: 33558762 , PubMed: 36170811). The composition of the serine palmitoyltransferase (SPT) complex determines the substrate preference (PubMed: 19416851 , PubMed: 33558762). The SPTLC1-SPTLC2-SPTSSA complex shows a strong preference for C16-CoA substrate, while the SPTLC1-SPTLC3-SPTSSA isozyme uses both C14-CoA and C16-CoA as substrates, with a slight preference for C14-CoA (PubMed: 19416851 , PubMed: 19648650). The SPTLC1-SPTLC2-SPTSSB complex shows a strong preference for C18-CoA substrate, while the SPTLC1-SPTLC3-SPTSSB isozyme displays an ability to use a broader range of acyl-CoAs, without apparent preference (PubMed: 19416851 , PubMed: 19648650 , PubMed: 33558761 , PubMed: 33558762). Required for adipocyte cell viability and metabolic homeostasis (By similarity).
Cellular Location	Endoplasmic reticulum membrane; Single-pass membrane protein {ECO:0000250 UniProtKB:O35704}
Tissue Location	Widely expressed. Not detected in small intestine.

Background

SPTLC1(Serine palmitoyltransferase, long chain base subunit 1), also known as SPT1, LCB1, is a protein which in humans is encoded by the SPTLC1 gene. Dawkins et al.(2001)noted that the SPTLC1 gene maps to chromosome 9q22.1-q22.3. Serine palmitoyltransferase, which consists of two different subunits, is the initial enzyme in sphingolipid biosynthesis. It converts L-serine and palmitoyl CoA to 3-oxosphinganine with pyridoxal 5'-phosphate as a cofactor. The product of this gene is the long chain base subunit 1 of serine palmitoyltransferase. Mutations in this gene were identified in patients with hereditary sensory neuropathy type 1. Alternatively spliced variants encoding different isoforms have been identified.

Images

Anti-SPTLC1 antibody, ABO11084, Western blottingAll lanes: Anti SPTLC1 (ABO11084) at 0.5ug/mlLane 1: SW620 Whole Cell Lysate at 40ugLane 2: S549 Whole Cell Lysate at 40ugLane 3: PANC Whole Cell Lysate at 40ugLane 4: U87 Whole Cell Lysate at 40ugPredicted bind size: 53KDObserved bind size: 53KD



Anti-SPTLC1 antibody, ABO11084, IHC(P)IHC(P): Human Mammary Cancer Tissue

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.