

Anti-Myoglobin Antibody

Catalog # ABO11134

Product Information

Application	WB
Primary Accession	P02144
Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for Myoglobin(MB) detection. Tested with WB in Human.
Reconstitution	Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	4151
Other Names	Myoglobin, MB
Calculated MW	17184
Application Details	Western blot, 0.1-0.5 µg/ml, Human
Source	Eukaryota
Protein Name	myoglobin
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg Thimerosal, 0.05mg Na ₃ .
Immunogen	A synthetic peptide corresponding to a sequence at the C-terminus of human Myoglobin(138-154aa LFRKDMASNYKELGFQG).
Purification	Immunogen affinity purified.
Cross Reactivity	No cross reactivity with other proteins
Storage	At -20 °C for one year. After reconstitution, at 4 °C for one month. It can also be aliquotted and stored frozen at -20 °C for a longer time. Avoid repeated freezing and thawing.
Sequence Similarities	Belongs to the globin family.

Protein Information

Name	MB (HGNC:6915)
Function	Monomeric heme protein which primary function is to store oxygen and facilitate its diffusion within muscle tissues. Reversibly binds oxygen through a pentacoordinated heme iron and enables its timely and efficient release as needed during periods of heightened demand (PubMed: 30918256 , PubMed: 34679218). Depending on the oxidative conditions of tissues and cells, and in addition to its ability to bind oxygen, it also has a nitrite reductase activity whereby it regulates the production of bioactive nitric oxide (PubMed: 32891753). Under stress conditions, like hypoxia and anoxia, it also protects cells against reactive oxygen species thanks to its pseudoperoxidase activity (PubMed: 34679218).
Cellular Location	Cytoplasm, sarcoplasm

Background

Myoglobin(MB) also known as PVALB, is a single-chain globular protein of 153 or 154 amino acids, containing a heme (iron-containing porphyrin) prosthetic group in the center around which the remaining apoprotein folds. It is a member of the globin superfamily and is expressed in skeletal and cardiac muscles. This gene is mapped to chromosome 22q11-q13. Myoglobin is released from damaged muscle tissue (rhabdomyolysis), which has very high concentrations of myoglobin. The released myoglobin is filtered by the kidneys but is toxic to the renal tubular epithelium and so may cause acute renal failure.

Images



Anti-Myoglobin antibody, ABO11134, Western blotting
Lane 1: Human Placenta Tissue Lysate
Lane 2: Human Placenta Tissue Lysate

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.