

Anti-RIP2 Antibody

Catalog # ABO11169

Product Information

Application	WB
Primary Accession	O43353
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for Receptor-interacting serine/threonine-protein kinase 2(RIPK2) detection. Tested with WB in Human;Mouse;Rat.
Reconstitution	Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	8767
Other Names	Receptor-interacting serine/threonine-protein kinase 2, 2.7.11.1, CARD-containing interleukin-1 beta-converting enzyme-associated kinase, CARD-containing IL-1 beta ICE-kinase, RIP-like-interacting CLARP kinase, Receptor-interacting protein 2, RIP-2, Tyrosine-protein kinase RIPK2, 2.7.10.2, RIPK2, CARDIAK, RICK, RIP2
Calculated MW	61195
Application Details	Western blot, 0.1-0.5 µg/ml, Human, Rat, Mouse
Subcellular Localization	Cytoplasm .
Tissue Specificity	Detected in heart, brain, placenta, lung, peripheral blood leukocytes, spleen, kidney, testis, prostate, pancreas and lymph node.
Source	Eukaryota
Protein Name	Receptor-interacting serine/threonine-protein kinase 2
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg Thimerosal, 0.05mg Na ₃ N.
Immunogen	A synthetic peptide corresponding to a sequence at the C-terminus of human RIP2(495-514aa DIQGEEFAKVIVQKLKDNKQ), different from the related rat and mouse sequences by one amino acid.
Purification	Immunogen affinity purified.

Cross Reactivity	No cross reactivity with other proteins
Storage	At -20 °C for one year. After r ° Constitution, at 4 °C for one month. It ° Can also be aliquotted and stored frozen at -20 °C for a longer time.Avoid repeated freezing and thawing.
Sequence Similarities	Belongs to the protein kinase superfamily. TKL Ser/Thr protein kinase family.

Protein Information


Name	RIPK2 {ECO:0000303 PubMed:30026309, ECO:0000312 HGNC:HGNC:10020}
Function	Serine/threonine/tyrosine-protein kinase that plays an essential role in modulation of innate and adaptive immune responses (PubMed:14638696, PubMed:17054981, PubMed:21123652, PubMed:28656966, PubMed:9575181, PubMed:9642260). Acts as a key effector of NOD1 and NOD2 signaling pathways: upon activation by bacterial peptidoglycans, NOD1 and NOD2 oligomerize and recruit RIPK2 via CARD-CARD domains, leading to the formation of RIPK2 filaments (PubMed:17054981, PubMed:17562858, PubMed:21123652, PubMed:22607974, PubMed:28656966, PubMed:29452636, PubMed:30026309). Once recruited, RIPK2 autophosphorylates and undergoes 'Lys-63'-linked polyubiquitination by E3 ubiquitin ligases XIAP, BIRC2 and BIRC3, as well as 'Met-1'-linked (linear) polyubiquitination by the LUBAC complex, becoming a scaffolding protein for downstream effectors (PubMed:22607974, PubMed:28545134, PubMed:29452636, PubMed:30026309, PubMed:30279485, PubMed:30478312). 'Met-1'-linked polyubiquitin chains attached to RIPK2 recruit IKKKG/NEMO, which undergoes 'Lys-63'-linked polyubiquitination in a RIPK2-dependent process (PubMed:17562858, PubMed:22607974, PubMed:29452636, PubMed:30026309). 'Lys-63'-linked polyubiquitin chains attached to RIPK2 serve as docking sites for TAB2 and TAB3 and mediate the recruitment of MAP3K7/TAK1 to IKKKG/NEMO, inducing subsequent activation of IKKKB/IKKB (PubMed:18079694). In turn, NF-kappa-B is released from NF-kappa-B inhibitors and translocates into the nucleus where it activates the transcription of hundreds of genes involved in immune response, growth control, or protection against apoptosis (PubMed:18079694). The protein kinase activity is dispensable for the NOD1 and NOD2 signaling pathways (PubMed:29452636, PubMed:30026309). Contributes to the tyrosine phosphorylation of the guanine exchange factor ARHGEF2 through Src tyrosine kinase leading to NF-kappa-B activation by NOD2 (PubMed:21887730). Also involved in adaptive immunity: plays a role during engagement of the T-cell receptor (TCR) in promoting BCL10 phosphorylation and subsequent NF-kappa-B activation (PubMed:14638696). Plays a role in the inactivation of RHOA in response to NGFR signaling (PubMed:26646181).
Cellular Location	Cytoplasm. Cell membrane; Peripheral membrane protein. Endoplasmic reticulum. Note=Recruited to the cell membrane by NOD2 following stimulation by bacterial peptidoglycans
Tissue Location	Detected in heart, brain, placenta, lung, peripheral blood leukocytes, spleen, kidney, testis, prostate, pancreas and lymph node.

Background

RIPK2(Receptor-interacting serine/threonine-protein kinase 2), also known as CARD3, CARDIAK, RICK, RIP2, is an enzyme that in humans is encoded by the RIPK2 gene. It has 540-amino acid protein in length. Northern blot analysis revealed that RICK is expressed in various human tissues as 2.5- and 1.8-kb

mRNAs that differ due to alternative polyadenylation. RICK is a novel kinase that may regulate apoptosis induced by the FAS receptor pathway. This gene encodes a member of the receptor-interacting protein(RIP) family of serine/threonine protein kinases. The encoded protein contains a C-terminal χ caspase recruitment domain(CARD), and is a component of signaling complexes in both the innate and adaptive immune pathways. It is a potent activator of χ NF-kappa B χ and inducer of apoptosis in response to various stimuli, CARDIAK(CARD-containing ICE-associated kinase) specifically interacted with the CARD of ICE/caspase-1, and this interaction correlated with the processing of pro-caspase-1 and the formation of the active caspase-1 p20 subunit.

Images



Anti-RIP2 antibody, ABO11169, Western blottingAll lanes:
 Anti RIP2 (ABO11169) at 0.5ug/mlLane 1: A549 Whole Cell
 Lysate at 40ugLane 2: HELA Whole Cell Lysate at
 40ugLane 3: PANC Whole Cell Lysate at 40ugLane 4:
 COLO320 Whole Cell Lysate at 40ugPredicted bind size:
 61KDObserved bind size: 61KD

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.