

Anti-Flt-3ligand Antibody

Catalog # ABO11217

Product Information

Application	WB
Primary Accession	P49771
Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for Fms-related tyrosine kinase 3 ligand(FLT3LG) detection. Tested with WB in Human.
Reconstitution	Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	2323
Other Names	Fms-related tyrosine kinase 3 ligand, Flt3 ligand, Flt3L, SL cytokine, FLT3LG
Calculated MW	26416
Application Details	Western blot, 0.1-0.5 µg/ml, Human
Subcellular Localization	Isoform 1: Cell membrane; Single-pass type I membrane protein.
Source	Eukaryota
Protein Name	Fms-related tyrosine kinase 3 ligand
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg Thimerosal, 0.05mg NaN3.
Immunogen	A synthetic peptide corresponding to a sequence at the C-terminus of human Flt-3ligand(206-220aa HWQRTRRRTPRPGEQ).
Purification	Immunogen affinity purified.
Cross Reactivity	No cross reactivity with other proteins
Storage	At -20 °C for one year. After reconstitution, at 4 °C for one month. It can also be aliquotted and stored frozen at -20 °C for a longer time. Avoid repeated freezing and thawing.


Protein Information

Name	FLT3LG
Function	Stimulates the proliferation of early hematopoietic cells by activating FLT3. Synergizes well with a number of other colony stimulating factors and interleukins. Required for the development of B cells, and dendritic cells (DCs).
Cellular Location	[Isoform 1]: Cell membrane; Single-pass type I membrane protein

Background

FLT3LG(FMS-Related Tyrosine Kinase 3 Ligand) also called FLT3 LIGAND, FL or FLT3L, is a protein which in humans is encoded by the FLT3LG gene. FLT3LG controls the development of DCs and is particularly important for plasmacytoid DCs and CD8-positive classical DCs and their CD103-positive tissue counterparts. Flt3 ligand(FL) is a hematopoietic four helical bundle cytokine. It is structurally homologous to stem cell factor(SCF) and colony stimulating factor 1(CSF-1). In synergy with other growth factors, Flt3 ligand stimulates the proliferation and differentiation of various blood cell progenitors. Lyman et al.(1993) found that Flt3 ligand stimulated proliferation of hematopoietic progenitor cells isolated from mouse fetal liver or adult mouse bone marrow. Hannum et al.(1994) concluded that FL enhances the response of stem and primitive progenitor cells to other growth factors to generate all myeloid lineages except erythroid cells. Assays of C-terminally truncated FL proteins confirmed that the N-terminal half conferred FL activity. Depletion of the Pi3k -Mtor negative regulator Pten facilitated Flt3l-driven DC development in culture.

Images



Anti-Flt-3ligand antibody, ABO11217, Western blotting
 Recombinant Protein Detection Source: E.coli derived -recombinant Human Flt-3ligand, 40.7KD(162aa tag+P36-H235)
 Lane 1: Recombinant Human Flt-3ligand Protein 10ng
 Lane 2: Recombinant Human Flt-3ligand Protein 5ng

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.