

Anti-mtTFA Antibody

Catalog # ABO11239

Product Information

Application	WB, IHC-P, IHC-F, ICC
Primary Accession	Q00059
Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for Transcription factor A, mitochondrial(TFAM) detection. Tested with WB, IHC-P, IHC-F, ICC in Human.
Reconstitution	Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	7019
Other Names	Transcription factor A, mitochondrial, mtTFA, Mitochondrial transcription factor 1, MtTF1, Transcription factor 6, TCF-6, Transcription factor 6-like 2, TFAM, TCF6, TCF6L2
Calculated MW	29097
Application Details	Immunohistochemistry(Paraffin-embedded Section), 0.5-1 μ g/ml, Human, By Heat Immunocytochemistry , 0.5-1 μ g/ml, Human, - Immunohistochemistry(Frozen Section), 0.5-1 μ g/ml, Human, - Western blot, 0.1-0.5 μ g/ml, Human
Subcellular Localization	Mitochondrion. Mitochondrion matrix, mitochondrion nucleoid.
Source	Eukaryota
Protein Name	Transcription factor A, mitochondrial
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg Thimerosal, 0.05mg Na ₃ .
Immunogen	A synthetic peptide corresponding to a sequence at the C-terminus of human mtTFA(227-246aa RKDLLRRTIKKQRKYGAEEC).
Purification	Immunogen affinity purified.
Cross Reactivity	No cross reactivity with other proteins

Storage	At -20° C for one year. After r° Constitution, at 4° C for one month. It° Can also be aliquotted and stored frozen at -20° C for a longer time.Avoid repeated freezing and thawing.
----------------	---

Protein Information



Name	TFAM (HGNC:11741)
Synonyms	TCF6, TCF6L2
Function	Binds to the mitochondrial light strand promoter and functions in mitochondrial transcription regulation (PubMed: 29445193 , PubMed: 32183942). Component of the mitochondrial transcription initiation complex, composed at least of TFB2M, TFAM and POLRMT that is required for basal transcription of mitochondrial DNA (PubMed: 29149603). In this complex, TFAM recruits POLRMT to a specific promoter whereas TFB2M induces structural changes in POLRMT to enable promoter opening and trapping of the DNA non-template strand (PubMed: 20410300). Required for accurate and efficient promoter recognition by the mitochondrial RNA polymerase (PubMed: 22037172). Promotes transcription initiation from the HSP1 and the light strand promoter by binding immediately upstream of transcriptional start sites (PubMed: 22037172). Is able to unwind DNA (PubMed: 22037172). Bends the mitochondrial light strand promoter DNA into a U-turn shape via its HMG boxes (PubMed: 1737790). Required for maintenance of normal levels of mitochondrial DNA (PubMed: 19304746 , PubMed: 22841477). May play a role in organizing and compacting mitochondrial DNA (PubMed: 22037171).
Cellular Location	Mitochondrion. Mitochondrion matrix, mitochondrion nucleoid

Background


TFAM(Transcription factor A, mitochondrial), also known as TCF6 or TCF6L2, is a 162-amino acid protein that activates transcription of each mitochondrial DNA(mtDNA) strand by binding to an element of approximately 30 nucleotides present in both the light-strand and the heavy-strand promoters. By Southern blot analysis of restriction enzyme digests of human/Chinese hamster somatic cell hybrid lines, Milatovich et al.(1992) mapped TFAM sequences, which they called MTTF1, to 3 different chromosomes: chromosomes 10, 7p, and 11q. By PCR-based screening of a somatic cell hybrid panel and by fluorescence in situ hybridization, Scott(2007) stated that the sequences mapped to chromosomes 7p(TCF6L1) and 11q(MTTF1, or TCF6L3) are pseudogenes. Larsson et al.(1997) mapped the mouse mitochondrial transcription factor A gene(Tfam) to the central part of mouse chromosome 10. This region exhibits syntenic homology with human 10q21. Mitochondrial transcription factor A is a key activator of mitochondrial transcription in mammals. It also has a role in mitochondrial DNA replication, since transcription generates an RNA primer necessary for initiation of mtDNA replication.

Images


Anti-mtTFA antibody, ABO11239, Western blotting
Lane 1: HELA Cell Lysate
Lane 2: JURKAT Cell Lysate
Lane 3: 293T Cell Lysate
Lane 4: Rabbit igG(55kD)
Lane 5: A431 Cell Lysate
Lane 6: RAJI Cell Lysate
Lane 7: CEM Cell Lysate
Lane 8: HL-60 Cell Lysate
Lane 9: HUT Cell Lysate



Anti-mtTFA antibody, ABO11239, IHC(P)IHC(P): Human Intestinal Cancer Tissue



Anti-mtTFA antibody, ABO11239, ICCICC: HELA Cell



Anti-mtTFA antibody, ABO11239, IHC(F)IHC(F): Human Placenta Tissue

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.