

Anti-TJP2 Antibody

Catalog # ABO11279

Product Information

Application	WB, IHC-P
Primary Accession	Q9Z0U1
Host	Rabbit
Reactivity	Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for Tight junction protein ZO-2(TJP2) detection. Tested with WB, IHC-P in Mouse;Rat.
Reconstitution	Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	21873
Other Names	Tight junction protein ZO-2, Tight junction protein 2, Zona occludens protein 2, Zonula occludens protein 2, Tjp2, Zo2
Calculated MW	131280
Application Details	Immunohistochemistry(Paraffin-embedded Section), 0.5-1 µg/ml, Mouse, Rat, By Heat Western blot, 0.1-0.5 µg/ml, Mouse, Rat
Subcellular Localization	Cell membrane ; Peripheral membrane protein ; Cytoplasmic side . Nucleus . Cell junction, tight junction . Also nuclear under environmental stress conditions and in migratory endothelial cells and subconfluent epithelial cell cultures. .
Source	Eukaryota
Protein Name	Tight junction protein ZO-2
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg Thimerosal, 0.05mg NaN3.
Immunogen	A synthetic peptide corresponding to a sequence at the C-terminus of mouse TJP2(1149-1167aa AHSKRGYYSQPSRYRDTL), identical to the related rat sequence.
Purification	Immunogen affinity purified.
Cross Reactivity	No cross reactivity with other proteins

Storage	At -20 °C for one year. After r ° Constitution, at 4 °C for one month. It ° Can also be aliquotted and stored frozen at -20 °C for a longer time.Avoid repeated freezing and thawing.
Sequence Similarities	Belongs to the MAGUK family.


Protein Information

Name	Tjp2 {ECO:0000312 MGI:MGI:1341872}
Function	Plays a role in tight junctions and adherens junctions (PubMed: 10026224). Acts as a positive regulator of RANKL-induced osteoclast differentiation, potentially via mediating downstream transcriptional activity (PubMed: 22437732).
Cellular Location	Cell junction, adherens junction. Cell membrane; Peripheral membrane protein; Cytoplasmic side. Nucleus. Cell junction, tight junction. Note=Also nuclear under environmental stress conditions and in migratory endothelial cells and subconfluent epithelial cell cultures. Localizes to tight junctions during initial stages of their formation (By similarity). {ECO:0000250, ECO:0000250 UniProtKB:Q95168}

Background


TJP2(Tight Junction Protein 2), also known as Zona Occludens 2 or ZO2, is a protein that in humans is encoded by the TJP2 gene. Tight junction proteins(TJPs) belong to a family of membrane-associated guanylate kinase(MAGUK) homologs that are involved in the organization of epithelial and endothelial intercellular junctions. Duclos et al.(1994) mapped the TJP2 gene telomeric to the Friedreich ataxia critical region on chromosome 9q13-q21. TJP2 lies about 70 kb centromeric to the X123 gene and is transcribed in the centromere-to-telomere direction. Using in vitro assays and immunoprecipitation studies, Itoh et al.(1999) showed that the mouse Tjp1, Tjp2, and Tjp3 PDZ1 domains interacted with the C-terminal cytoplasmic domains of Cldn1 through Cldn8. In the mouse inner ear, Walsh et al.(2010) found that Tjp2 expression decreased rapidly between E16.5 and age 1 week to a level in adult mice that was approximately 50% of the level at birth(P0).

Images



Anti-TJP2 antibody, ABO11279, Western blotting
Lane 1: Rat Brain Tissue Lysate
Lane 2: PC-12 Cell Lysate
Lane 3: Mouse Brain Tissue Lysate
Lane 4: HEPA Cell Lysate

Anti-TJP2 antibody, ABO11279, IHC(P)
IHC(P): Mouse Liver Tissue



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.