

Anti-TORC1 Antibody

Catalog # ABO11295

Product Information

Application	WB
Primary Accession	Q6UUV9
Host	Rabbit
Reactivity	Human, Rat
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for CREB-regulated transcription coactivator 1(CRTC1) detection. Tested with WB in Human;Rat.
Reconstitution	Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	23373
Other Names	CREB-regulated transcription coactivator 1, Mucoepidermoid carcinoma translocated protein 1, Transducer of regulated cAMP response element-binding protein 1, TORC-1, Transducer of CREB protein 1, CRTC1 (HGNC:16062)
Calculated MW	67300
Application Details	Western blot, 0.1-0.5 µg/ml, Human, Rat
Subcellular Localization	Cytoplasm. Nucleus. Cytoplasmic when phosphorylated by SIK or AMPK and when sequestered by 14-3-3 proteins (By similarity). Translocated to the nucleus on Ser-151 dephosphorylation, instigated by a number of factors including calcium ion and cAMP levels. Light stimulation triggers a nuclear accumulation in the suprachiasmatic nucleus (SCN) of the brain. .
Tissue Specificity	Highly expressed in adult and fetal brain. Located to specific regions such as the prefrontal cortex and cerebellum. Very low expression in other tissues such as heart, spleen, lung, skeletal muscle, salivary gland, ovary and kidney. .
Source	Eukaryota
Protein Name	CREB-regulated transcription coactivator 1
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg Thimerosal, 0.05mg NaN3.
Immunogen	A synthetic peptide corresponding to a sequence at the C-terminus of human TORC1(506-521aa EQQMAARQANALSHQL), different from the related rat and

mouse sequences by two amino acids.

Purification	Immunogen affinity purified.
Cross Reactivity	No cross reactivity with other proteins
Storage	At -20 °C for one year. After r ° Constitution, at 4 °C for one month. It ° Can also be aliquotted and stored frozen at -20 °C for a longer time.Avoid repeated freezing and thawing.
Sequence Similarities	Belongs to the TORC family.


Protein Information

Name	CRTC1 (HGNC:16062)
Function	Transcriptional coactivator for CREB1 which activates transcription through both consensus and variant cAMP response element (CRE) sites. Acts as a coactivator, in the SIK/TORC signaling pathway, being active when dephosphorylated and acts independently of CREB1 'Ser-133' phosphorylation. Enhances the interaction of CREB1 with TAF4. Regulates the expression of specific CREB-activated genes such as the steroidogenic gene, StAR. Potent coactivator of PGC1alpha and inducer of mitochondrial biogenesis in muscle cells. In the hippocampus, involved in late-phase long-term potentiation (L-LTP) maintenance at the Schaffer collateral-CA1 synapses. May be required for dendritic growth of developing cortical neurons (By similarity). In concert with SIK1, regulates the light-induced entrainment of the circadian clock. In response to light stimulus, coactivates the CREB-mediated transcription of PER1 which plays an important role in the photic entrainment of the circadian clock.
Cellular Location	Cytoplasm. Nucleus. Note=Cytoplasmic when phosphorylated by SIK or AMPK and when sequestered by 14-3-3 proteins (PubMed:16817901) Translocated to the nucleus on Ser-151 dephosphorylation, instigated by a number of factors including calcium ion and cAMP levels (PubMed:15589160). Light stimulation triggers a nuclear accumulation in the suprachiasmatic nucleus (SCN) of the brain (By similarity) {ECO:0000250 UniProtKB:Q68ED7, ECO:0000269 PubMed:15589160, ECO:0000269 PubMed:16817901}
Tissue Location	Highly expressed in adult and fetal brain. Located to specific regions such as the prefrontal cortex and cerebellum. Very low expression in other tissues such as heart, spleen, lung, skeletal muscle, salivary gland, ovary and kidney.

Background

CRTC1(CREB-Regulated Transcription Coactivator 1), also known as MECT1, TORC1 or KIAA0616, is a protein that in humans is encoded by the CRTC1 gene. By sequence analysis, Tonon et al.(2003) mapped the CRTC1 gene to chromosome 19p13. Kovacs et al.(2007) found that Torc1 was expressed in adult mouse brain and cultured neurons, and that it translocated to the nucleus upon concomitant activation of calcium and cAMP signaling pathways. Mair et al.(2011) demonstrated that CRTC1 is a direct AMPK target, and interacts with the CREB homolog-1(CRH1) transcription factor in vivo. The prolongevity effects of activating AMPK or deactivating calcineurin decrease CRTC1 and CRH1 activity and induce transcriptional responses similar to those of CRH1-null worms.

Images



Anti-TORC1 antibody, ABO11295, Western blotting
 Recombinant Protein Detection Source: E.coli derived -recombinant Human CRTC1, 29.5KD(162aa tag+ L460-Y568)
 Lane 1: Recombinant Human CRTC1 Protein 10ng
 Lane 2: Recombinant Human CRTC1 Protein 5ng
 Lane 3: Recombinant Human CRTC1 Protein 2.5ng

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.