

Anti-TrkC Antibody

Catalog # ABO11300

Product Information

Application	WB
Primary Accession	Q16288
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for NT-3 growth factor receptor(NTRK3) detection. Tested with WB in Human;Mouse;Rat.
Reconstitution	Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	4916
Other Names	NT-3 growth factor receptor, 2.7.10.1, GP145-TrkC, Trk-C, Neurotrophic tyrosine kinase receptor type 3, TrkC tyrosine kinase, NTRK3, TRKC
Calculated MW	94428
Application Details	Western blot, 0.1-0.5 µg/ml, Human, Mouse, Rat
Subcellular Localization	Membrane; Single-pass type I membrane protein.
Tissue Specificity	Widely expressed but mainly in nervous tissue. Isoform 2 is expressed at higher levels in adult brain than in fetal brain.
Source	Eukaryota
Protein Name	NT-3 growth factor receptor
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg Thimerosal, 0.05mg Na ₃ N.
Immunogen	A synthetic peptide corresponding to a sequence at the N-terminus of human TrkC(172-186aa QLWQEQGEAKLNSQN), different from the related rat and mouse sequences by three amino acids.
Purification	Immunogen affinity purified.
Cross Reactivity	No cross reactivity with other proteins
Storage	At -20 °C for one year. After reconstitution, at 4 °C for one month. It can also

Sequence Similarities

be aliquotted and stored frozen at -20 °C for a longer time. Avoid repeated freezing and thawing.
Belongs to the protein kinase superfamily. Tyr protein kinase family. Insulin receptor subfamily.


Protein Information

Name	NTRK3
Synonyms	TRKC
Function	Receptor tyrosine kinase involved in nervous system and probably heart development. Upon binding of its ligand NTF3/neurotrophin-3, NTRK3 autophosphorylates and activates different signaling pathways, including the phosphatidylinositol 3-kinase/AKT and the MAPK pathways, that control cell survival and differentiation.
Cellular Location	Membrane; Single-pass type I membrane protein.
Tissue Location	Widely expressed but mainly in nervous tissue. Isoform 2 is expressed at higher levels in adult brain than in fetal brain

Background

NTRK3 (Neurotrophic Tyrosine Kinase Receptor Type 3), also known as TRKC, is a protein that in humans is encoded by the NTRK3 gene. By PCR analysis of a somatic cell hybrid panel and by fluorescence in situ hybridization with the cDNA clone, McGregor et al. (1994) mapped the NTRK3 gene to 15q24-q25. Lamballe et al. (1991) isolated and characterized TRKC, a member of the TRK family of tyrosine protein kinase genes. They found that TRKC is preferentially expressed in the brain; in situ hybridization studies showed transcripts in the hippocampus, cerebral cortex, and the granular cell layer of the cerebellum. By functional studies in HeLa cells, Muinos-Gimeno et al. (2009) demonstrated that 5 miRNAs regulate the truncated form of NTRK3.

Images



Anti-TrkC antibody, ABO11300, Western blotting
Lane 1: Rat Brain Tissue Lysate
Lane 2: Mouse Brain Tissue Lysate
Lane 3: U87 Cell Lysate

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.