

Anti-T-bet/Tbx21 Antibody

Catalog # ABO11308

Product Information

Application	WB
Primary Accession	Q9UL17
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for T-box transcription factor TBX21(TBX21) detection. Tested with WB in Human;Mouse;Rat.
Reconstitution	Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	30009
Other Names	T-box transcription factor TBX21, T-box protein 21, T-cell-specific T-box transcription factor T-bet, Transcription factor TBLYM, TBX21, TBET, TBLYM
Calculated MW	58328
Application Details	Western blot, 0.1-0.5 µg/ml, Human, Rat, Mouse
Subcellular Localization	Nucleus.
Tissue Specificity	T-cell specific.
Source	Eukaryota
Protein Name	T-box transcription factor TBX21
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg Thimerosal, 0.05mg NaN ₃ .
Immunogen	A synthetic peptide corresponding to a sequence at the C-terminus of human T-bet(522-535aa DKEAEGQFYNYFPN), different from the related rat and mouse sequences by one amino acid.
Purification	Immunogen affinity purified.
Cross Reactivity	No cross reactivity with other proteins
Storage	At -20 °C for one year. After reconstitution, at 4 °C for one month. It can also be aliquotted and stored frozen at -20 °C for a longer time. Avoid repeated

freezing and thawing.

Protein Information


Name	TBX21
Synonyms	TBET, TBLYM
Function	Lineage-defining transcription factor which initiates Th1 lineage development from naive Th precursor cells both by activating Th1 genetic programs and by repressing the opposing Th2 and Th17 genetic programs (PubMed:10761931). Activates transcription of a set of genes important for Th1 cell function, including those encoding IFN- gamma and the chemokine receptor CXCR3. Induces permissive chromatin accessibility and CpG methylation in IFNG (PubMed:33296702). Activates IFNG and CXCR3 genes in part by recruiting chromatin remodeling complexes including KDM6B, a SMARCA4-containing SWI/SNF-complex, and an H3K4me2-methyltransferase complex to their promoters and all of these complexes serve to establish a more permissive chromatin state conducive with transcriptional activation (By similarity). Can activate Th1 genes also via recruitment of Mediator complex and P-TEFb (composed of CDK9 and CCNT1/cyclin-T1) in the form of the super elongation complex (SEC) to super-enhancers and associated genes in activated Th1 cells (PubMed:27292648). Inhibits the Th17 cell lineage commitment by blocking RUNX1-mediated transactivation of Th17 cell-specific transcriptinal regulator RORC. Inhibits the Th2 cell lineage commitment by suppressing the production of Th2 cytokines, such as IL-4, IL-5, and IL- 13, via repression of transcriptional regulators GATA3 and NFATC2. Protects Th1 cells from amplifying aberrant type-I IFN response in an IFN-gamma abundant microenvironment by acting as a repressor of type-I IFN transcription factors and type-I IFN-stimulated genes. Acts as a regulator of antiviral B-cell responses; controls chronic viral infection by promoting the antiviral antibody IgG2a isotype switching and via regulation of a broad antiviral gene expression program (By similarity). Required for the correct development of natural killer (NK) and mucosal-associated invariant T (MAIT) cells (PubMed:33296702).
Cellular Location	Nucleus
Tissue Location	T-cell specific..

Background

TBX21(T-Box 21), also called TBET, is a protein that in humans is encoded by the TBX21 gene. This gene is a member of a phylogenetically conserved family of genes that share a common DNA-binding domain, the T-box. Zhang and Yang(2000) mapped the mouse Tbx21 gene to chromosome 11D in an area showing homology of synteny with human chromosome 17. Szabo et al.(2000) showed that TBX21 expression correlates with IFNG expression in Th1 and natural killer(NK) cells. Ectopic expression of TBX21 both transactivated the IFNG gene and induced endogenous IFNG production. Finotto et al.(2002) observed reduced expression of the T(H)1 transcription factor Tbet in T cells from airways of patients with asthma compared with that in T cells from airways of nonasthmatic patients, suggesting that loss of Tbet might be associated with asthma.

Images

Anti-T-bet/Tbx21 antibody, ABO11308, Western blotting
Recombinant Protein Detection Source: E.coli



derived -recombinant Human TBX21,34.7KD(162aa tag+
V381-N535)Lane 1: Recombinant Human TBX21 Protein
5ngLane 2: Recombinant Human TBX21 Protein
2.5ngLane 3: Recombinant Human TBX21 Protein 1.25ng

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.