

Anti-Eif3b Antibody

Catalog # ABO11337

Product Information

Application	WB
Primary Accession	P55884
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for Eukaryotic translation initiation factor 3 subunit B(EIF3B) detection. Tested with WB in Human;Mouse;Rat.
Reconstitution	Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	8662
Other Names	Eukaryotic translation initiation factor 3 subunit B {ECO:0000255 HAMAP-Rule:MF_03001}, eIF3b {ECO:0000255 HAMAP-Rule:MF_03001}, Eukaryotic translation initiation factor 3 subunit 9 {ECO:0000255 HAMAP-Rule:MF_03001}, Prt1 homolog, hPrt1, eIF-3-eta {ECO:0000255 HAMAP-Rule:MF_03001}, eIF3 p110 {ECO:0000255 HAMAP-Rule:MF_03001}, eIF3 p116, EIF3B {ECO:0000255 HAMAP-Rule:MF_03001}
Calculated MW	92482
Application Details	Western blot, 0.1-0.5 µg/ml, Human, Rat, Mouse
Subcellular Localization	Cytoplasm .
Source	Eukaryota
Protein Name	Eukaryotic translation initiation factor 3 subunit B
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg Thimerosal, 0.05mg Na ₃ .
Immunogen	A synthetic peptide corresponding to a sequence at the C-terminus of human eIF3B(755-775aa EDFRKYRKMAQELYMEQKNER), different from the related rat and mouse sequences by one amino acid.
Purification	Immunogen affinity purified.
Cross Reactivity	No cross reactivity with other proteins

Storage	At -20° C for one year. After r° Constitution, at 4° C for one month. It° Can also be aliquotted and stored frozen at -20° C for a longer time.Avoid repeated freezing and thawing.
Sequence Similarities	Belongs to the eIF-3 subunit B family.


Protein Information

Name	EIF3B {ECO:0000255 HAMAP-Rule:MF_03001}
Function	RNA-binding component of the eukaryotic translation initiation factor 3 (eIF-3) complex, which is required for several steps in the initiation of protein synthesis (PubMed: 17581632 , PubMed: 25849773 , PubMed: 27462815 , PubMed: 9388245). The eIF-3 complex associates with the 40S ribosome and facilitates the recruitment of eIF-1, eIF-1A, eIF-2:GTP:methionyl-tRNAi and eIF-5 to form the 43S pre- initiation complex (43S PIC). The eIF-3 complex stimulates mRNA recruitment to the 43S PIC and scanning of the mRNA for AUG recognition. The eIF-3 complex is also required for disassembly and recycling of post-termination ribosomal complexes and subsequently prevents premature joining of the 40S and 60S ribosomal subunits prior to initiation (PubMed: 17581632 , PubMed: 9388245). The eIF-3 complex specifically targets and initiates translation of a subset of mRNAs involved in cell proliferation, including cell cycling, differentiation and apoptosis, and uses different modes of RNA stem-loop binding to exert either translational activation or repression (PubMed: 25849773).
Cellular Location	Cytoplasm {ECO:0000255 HAMAP-Rule:MF_03001}. Cytoplasm, Stress granule. Note=Localizes to stress granules following cellular stress

Background

EIF3B(Eukaryotic Translation Initiation Factor 3, Subunit B), also called PRT1 or EIF3-p116, is a protein that in humans is encoded by the EIF3B gene. By searching an EST database, Methot et al.(1997) identified a cDNA encoding a human PRT1 homolog. Asano et al.(1997) demonstrated that the 115-kD component of HeLa cell eIF3 is actually composed of 2 proteins, p116(PRT1) and an unrelated 110-kD protein. Chaudhuri et al.(1997) isolated cDNAs encoding PRT1, which they called p110 based on the size of the corresponding component of rabbit eIF3.

Images



Anti-Eif3b antibody, ABO11337, Western blotting
Lane 1: HELA Cell Lysate
Lane 2: 293T Cell Lysate
Lane 3: A431 Cell Lysate

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.